

177th Eigenvector

$$N_e = 5 \quad s = \frac{1}{2} \quad m_s = -\frac{1}{2}$$

Irred. Representation : $\Gamma_{4,2}$

$$E_{177} = \frac{1}{3} \left(-J - 3t + 5U + 50W - 2 \cos(\theta_5) \sqrt{A_6} \right)$$

$$\begin{aligned} |\Psi_{177}\rangle &= |5, \frac{1}{2}, -\frac{1}{2}, \Gamma_{4,2}\rangle \\ &= C_{177,1} (|022d\rangle - |20d2\rangle - |2d02\rangle + |d220\rangle) \\ &+ C_{177,2} (|02d2\rangle + |0d22\rangle - |202d\rangle - |220d\rangle + |22d0\rangle + |2d20\rangle - |d022\rangle - |d202\rangle) \\ &+ C_{177,3} (|2dud\rangle - |2udd\rangle + |d2du\rangle + |dd2u\rangle + |ddu2\rangle - |dud2\rangle - |u2dd\rangle - |ud2d\rangle) \end{aligned}$$

$$\begin{aligned} C_{177-1} &= \frac{5t^2}{2} + 3Ut + 34Wt - U^2 + J(-t + U + 8W) \\ &+ \left(\frac{1}{18} \left(-A_{22}^2 + 3(J + 4t - 3U - 34W)A_{22} + 36(17t - 13U - 72W)W \right) \right) \\ C_{177-2} &= \frac{1}{3}t \left(J + 6t + U - 2W - \cos(\theta_5) \sqrt{A_6} \right) \\ C_{177-3} &= \frac{1}{6}t \left(-J + 12t - U + 2W - 2 \cos(\theta_5) \sqrt{A_6} \right) \\ N_{177} &= 2\sqrt{C_{177,1}^2 + 2(C_{177,2}^2 + C_{177,3}^2)} \end{aligned}$$