

# Hilbert complexes with mixed boundary conditions—Part 2: Elasticity complex

Dirk Pauly<sup>1</sup>  | Michael Schomburg<sup>2</sup>

<sup>1</sup>Institut für Analysis, Fakultät  
Mathematik, Technische Universität  
Dresden, Dresden, Germany

<sup>2</sup>Fakultät für Mathematik, Universität  
Duisburg-Essen, Essen, Germany

**Correspondence**

Dirk Pauly, Institut für Analysis, Fakultät  
Mathematik, Technische Universität  
Dresden, Dresden, Germany.  
Email: dirk.pauly@tu-dresden.de

Communicated by: M. Waurick

We show that the elasticity Hilbert complex with mixed boundary conditions on bounded strong Lipschitz domains is closed and compact. The crucial results are compact embeddings which follow by abstract arguments using functional analysis together with particular regular decompositions. Higher Sobolev order results are also proved. This paper extends recent results on the de Rham Hilbert complex with mixed boundary conditions from Pauly and Schomburg (2021, 2022) and recent results on the elasticity Hilbert complex with empty or full boundary conditions from Pauly and Zulehner (2020, 2022).

**KEYWORDS**

regular potentials, regular decompositions, compact embeddings, Hilbert complexes, mixed boundary conditions, elasticity complex

**MSC CLASSIFICATION**

58Axx, 58Jxx, 35A23, 35Q61

## 1 | INTRODUCTION

In this paper, we prove regular decompositions and resulting compact embeddings for the *elasticity complex*

$$\dots \xrightarrow{\quad} L^2(\Omega) \xrightarrow{\text{symGrad}} L^2_{\mathbb{S}}(\Omega) \xrightarrow{\text{RotRot}_{\mathbb{S}}^T} L^2_{\mathbb{S}}(\Omega) \xrightarrow{\text{Div}_{\mathbb{S}}} L^2(\Omega) \xrightarrow{\quad} \dots$$

This extends the corresponding results from Pauly and Schomburg<sup>1,2</sup> for the de Rham complex

$$\dots \xrightarrow{\quad} L^{q-1,2}(\Omega) \xrightarrow{d^{q-1}} L^{q,2}(\Omega) \xrightarrow{d^q} L^{q+1,2}(\Omega) \xrightarrow{\quad} \dots,$$

whose 3D version for vector proxies is given by

$$\dots \xrightarrow{\quad} L^2(\Omega) \xrightarrow{d^0 \hat{=} \text{grad}} L^2(\Omega) \xrightarrow{d^1 \hat{=} \text{rot}} L^2(\Omega) \xrightarrow{d^2 \hat{=} \text{div}} L^2(\Omega) \xrightarrow{\quad} \dots$$

We shall consider mixed boundary conditions on a bounded strong Lipschitz domain  $\Omega \subset \mathbb{R}^3$ .

Like the de Rham complex, the elasticity complex has the geometric structure of a *Hilbert complex*, that is,

$$\dots \xrightarrow{\quad} H_0 \xrightarrow{A_0} H_1 \xrightarrow{A_1} H_2 \xrightarrow{\quad} \dots, \quad R(A_0) \subset N(A_1),$$

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Mathematical Methods in the Applied Sciences* published by John Wiley & Sons Ltd.

where  $A_0$  and  $A_1$  are densely defined and closed (unbounded) linear operators between Hilbert spaces  $H_\ell$ . The corresponding *domain Hilbert complex* is denoted by

$$\cdots \xrightarrow{\cdots} D(A_0) \xrightarrow{A_0} D(A_1) \xrightarrow{A_1} H_2 \xrightarrow{\cdots} \cdots$$

In fact, we show that the assumptions of lemma 2.22<sup>1,2</sup> hold, which provides an elegant, abstract, and short way to prove the crucial compact embeddings

$$D(A_1) \cap D(A_0^*) \hookrightarrow H_1 \quad (1)$$

for the elasticity Hilbert complex. In principle, our general technique—compact embeddings by regular decompositions and Rellich's selection theorem—works for all Hilbert complexes known in the literature; see, for example, Arnold and Hu<sup>3</sup> for a comprehensive list of such Hilbert complexes.

Roughly speaking, a regular decomposition has the form

$$D(A_1) = H_1^+ + A_0 H_0^+,$$

with regular subspaces  $H_0^+ \subset D(A_0)$  and  $H_1^+ \subset D(A_1)$  such that the embeddings  $H_0^+ \hookrightarrow H_0$  and  $H_1^+ \hookrightarrow H_1$  are compact. The compactness is typically and simply given by Rellich's selection theorem, which justifies the notion “regular.” By applying  $A_1$ , any regular decomposition implies regular potentials

$$R(A_1) = A_1 H_1^+$$

by the complex property. The respective regular potential and decomposition operators

$$\mathcal{P}_{A_1} : R(A_1) \rightarrow H_1^+, \quad \mathcal{Q}_{A_1}^1 : D(A_1) \rightarrow H_1^+, \quad \mathcal{Q}_{A_1}^0 : D(A_1) \rightarrow H_0^+$$

are bounded and satisfy  $A_1 \mathcal{P}_{A_1} = \text{id}_{R(A_1)}$  as well as  $\text{id}_{D(A_1)} = \mathcal{Q}_{A_1}^1 + A_0 \mathcal{Q}_{A_1}^0$ .

Note that (1) implies several important results related to the particular Hilbert complex by the so-called FA-ToolBox, such as closed ranges, Friedrichs/Poincaré-type estimates, Helmholtz-type decompositions, and comprehensive solution theories; compare previous works<sup>4–7</sup> and references.<sup>8–11</sup>

For a historical overview on the compact embeddings (1) corresponding to the de Rham complex and Maxwell's equations, that is, Weck's or Weber–Weck–Picard's selection theorem, see, for example, the introductions in previous references,<sup>12,13</sup> the original papers,<sup>14–19</sup> and the recent state-of-the-art results for mixed boundary conditions and bounded weak Lipschitz domains in previous works.<sup>12,20,21</sup> Compact embeddings (1) corresponding to the biharmonic and the elasticity complex are given in references<sup>10,11</sup> and previous works,<sup>8,9</sup> respectively. Note that in the recent paper<sup>3</sup> similar results have been shown for no or full boundary conditions using an alternative and more algebraic approach, the so-called Bernstein–Gelfand–Gelfand resolution (BGG).

## 2 | ELASTICITY COMPLEXES I

Throughout this paper, let  $\Omega \subset \mathbb{R}^3$  be a *bounded strong Lipschitz domain* with boundary  $\Gamma$ , decomposed into two parts  $\Gamma_t$  and  $\Gamma_n := \Gamma \setminus \Gamma_t$  with some *relatively open and strong Lipschitz boundary part*  $\Gamma_t \subset \Gamma$ .

### 2.1 | Notations and preliminaries

We will strongly use the notations and results from our corresponding papers for the elasticity complex<sup>10,11</sup> and for the de Rham complex.<sup>1,2</sup> In particular, we recall sections 2 and 3<sup>1,2</sup> including the notion of *extendable domains*.

We utilize the standard Sobolev spaces from previous works,<sup>1,2</sup> for example, the usual Lebesgue and Sobolev spaces (scalar or tensor valued)  $L^2(\Omega)$  and  $H^k(\Omega)$  with  $k \in \mathbb{N}_0$ . Boundary conditions are introduced in the *strong sense* as closures of respective test fields, that is,

$$H_{\Gamma_t}^k(\Omega) := \overline{C_{\Gamma_t}^\infty(\Omega)}^{H^k(\Omega)},$$

we well as in the *weak sense* by

$$\mathbf{H}_{\Gamma_t}^k(\Omega) := \left\{ u \in H^k(\Omega) : \langle \partial^\alpha u, \phi \rangle_{L^2(\Omega)} = (-1)^{|\alpha|} \langle u, \partial^\alpha \phi \rangle_{L^2(\Omega)} \quad \forall \phi \in C_{\Gamma_n}^\infty(\Omega) \quad \forall |\alpha| \leq k \right\}.$$

**Lemma 2.1** (lemma 3.2 and theorem 4.6<sup>1,2</sup>).  $\mathbf{H}_{\Gamma_t}^k(\Omega) = H_{\Gamma_t}^k(\Omega)$ ; that is, weak and strong boundary conditions coincide for the standard Sobolev spaces with mixed boundary conditions.

We shall use the abbreviations  $\mathbf{H}_{\emptyset}^k(\Omega) = H^k(\Omega)$  and  $\mathbf{H}_{\Gamma_t}^0(\Omega) = L^2(\Omega)$ , where the first notion is actually a density result and incorporated into the notation by purpose.

## 2.2 | Operators

Let  $\text{symGrad}$ ,  $\text{RotRot}^\top$ , and  $\text{Div}$  (here  $\text{Grad}$ ,  $\text{Rot}$ , and  $\text{Div}$  act row-wise as the operators  $\text{grad}$ ,  $\text{rot}$ , and  $\text{div}$  from the vector de Rham complex) be realized as densely defined (unbounded) linear operators

$$\begin{aligned} \text{sym}\mathring{\text{Grad}}_{\Gamma_t} : D(\text{sym}\mathring{\text{Grad}}_{\Gamma_t}) \subset L^2(\Omega) &\rightarrow L_{\mathbb{S}}^2(\Omega); & v &\mapsto \text{symGrad} v = \frac{1}{2} (\text{Grad} v + (\text{Grad} v)^\top), \\ \text{Rot}\mathring{\text{Rot}}_{\mathbb{S},\Gamma_t}^\top : D(\text{Rot}\mathring{\text{Rot}}_{\mathbb{S},\Gamma_t}^\top) \subset L_{\mathbb{S}}^2(\Omega) &\rightarrow L_{\mathbb{S}}^2(\Omega); & S &\mapsto \text{RotRot}^\top S = \text{Rot}((\text{Rot} S)^\top), \\ \mathring{\text{Div}}_{\mathbb{S},\Gamma_t} : D(\mathring{\text{Div}}_{\mathbb{S},\Gamma_t}) \subset L_{\mathbb{S}}^2(\Omega) &\rightarrow L^2(\Omega); & T &\mapsto \text{Div} T \end{aligned}$$

( $S, T, \text{Grad} v, \text{symGrad} v, \text{Rot} S, \text{RotRot}^\top S$  are  $(3 \times 3)$ -tensor fields, and  $v, \text{Div} T$  are 3-vector fields) with domains of definition

$$D(\text{sym}\mathring{\text{Grad}}_{\Gamma_t}) := C_{\Gamma_t}^\infty(\Omega), \quad D(\text{Rot}\mathring{\text{Rot}}_{\mathbb{S},\Gamma_t}^\top) := C_{\mathbb{S},\Gamma_t}^\infty(\Omega), \quad D(\mathring{\text{Div}}_{\mathbb{S},\Gamma_t}) := C_{\mathbb{S},\Gamma_t}^\infty(\Omega)$$

satisfying the complex properties

$$\text{Rot}\mathring{\text{Rot}}_{\mathbb{S},\Gamma_t}^\top \text{sym}\mathring{\text{Grad}}_{\Gamma_t} \subset 0, \quad \mathring{\text{Div}}_{\mathbb{S},\Gamma_t} \text{Rot}\mathring{\text{Rot}}_{\mathbb{S},\Gamma_t}^\top \subset 0.$$

For elementary properties of these operators, see, for example, references<sup>10,11</sup>; in particular, we have a collection of formulas presented in Lemma A.1. Here, we introduce the Lebesgue Hilbert space and the test space of symmetric tensor fields

$$L_{\mathbb{S}}^2(\Omega) := \{ S \in L^2(\Omega) : S^\top = S \}, \quad C_{\mathbb{S},\Gamma_t}^\infty(\Omega) := C_{\Gamma_t}^\infty(\Omega) \cap L_{\mathbb{S}}^2(\Omega),$$

respectively. We get the elasticity complex on smooth tensor fields

$$\dots \xrightarrow{\dots} L^2(\Omega) \xrightarrow{\text{sym}\mathring{\text{Grad}}_{\Gamma_t}} L_{\mathbb{S}}^2(\Omega) \xrightarrow{\text{Rot}\mathring{\text{Rot}}_{\mathbb{S},\Gamma_t}^\top} L_{\mathbb{S}}^2(\Omega) \xrightarrow{\mathring{\text{Div}}_{\mathbb{S},\Gamma_t}} L^2(\Omega) \xrightarrow{\dots} \dots$$

For a more structured introduction of the latter operators, see Appendix B1.

The closures

$$\text{symGrad}_{\Gamma_t} := \overline{\text{sym}\mathring{\text{Grad}}_{\Gamma_t}}, \quad \text{RotRot}_{\mathbb{S},\Gamma_t}^\top := \overline{\text{Rot}\mathring{\text{Rot}}_{\mathbb{S},\Gamma_t}^\top}, \quad \text{Div}_{\mathbb{S},\Gamma_t} := \overline{\mathring{\text{Div}}_{\mathbb{S},\Gamma_t}}$$

and Hilbert space adjoints

$$\text{symGrad}_{\Gamma_t}^* = \text{sym}\mathring{\text{Grad}}_{\Gamma_t}^*, \quad (\text{RotRot}_{\mathbb{S},\Gamma_t}^\top)^* = (\text{Rot}\mathring{\text{Rot}}_{\mathbb{S},\Gamma_t}^\top)^*, \quad \text{Div}_{\mathbb{S},\Gamma_t}^* = \mathring{\text{Div}}_{\mathbb{S},\Gamma_t}^*$$

are given by the densely defined and closed linear operators

$$\begin{aligned} A_0 &:= \text{symGrad}_{\Gamma_t} : D(\text{symGrad}_{\Gamma_t}) \subset L^2(\Omega) \rightarrow L_{\mathbb{S}}^2(\Omega); & v &\mapsto \text{symGrad} v, \\ A_1 &:= \text{RotRot}_{\mathbb{S},\Gamma_t}^\top : D(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top) \subset L_{\mathbb{S}}^2(\Omega) \rightarrow L_{\mathbb{S}}^2(\Omega); & S &\mapsto \text{RotRot}^\top S, \\ A_2 &:= \text{Div}_{\mathbb{S},\Gamma_t} : D(\text{Div}_{\mathbb{S},\Gamma_t}) \subset L_{\mathbb{S}}^2(\Omega) \rightarrow L^2(\Omega); & T &\mapsto \text{Div} T, \\ A_0^* &= \text{symGrad}_{\Gamma_t}^* = -\mathring{\text{Div}}_{\mathbb{S},\Gamma_n} : D(\mathring{\text{Div}}_{\mathbb{S},\Gamma_n}) \subset L_{\mathbb{S}}^2(\Omega) \rightarrow L^2(\Omega); & S &\mapsto -\text{Div} S, \\ A_1^* &= (\text{Rot}\mathring{\text{Rot}}_{\mathbb{S},\Gamma_t}^\top)^* = \mathbf{Rot}\mathbf{Rot}_{\mathbb{S},\Gamma_n}^\top : D(\mathbf{Rot}\mathbf{Rot}_{\mathbb{S},\Gamma_n}^\top) \subset L_{\mathbb{S}}^2(\Omega) \rightarrow L_{\mathbb{S}}^2(\Omega); & T &\mapsto \text{RotRot}^\top T, \\ A_2^* &= \text{Div}_{\mathbb{S},\Gamma_t}^* = -\mathbf{sym}\mathbf{Grad}_{\Gamma_n} : D(\mathbf{sym}\mathbf{Grad}_{\Gamma_n}) \subset L^2(\Omega) \rightarrow L_{\mathbb{S}}^2(\Omega); & v &\mapsto -\text{symGrad} v \end{aligned}$$

with domains of definition

$$\begin{aligned} D(A_0) &= D(\text{symGrad}_{\Gamma_t}) = \mathbf{H}_{\Gamma_t}(\text{symGrad}, \Omega), & D(A_0^*) &= D(\mathbf{Div}_{\mathbb{S}, \Gamma_n}) = \mathbf{H}_{\mathbb{S}, \Gamma_n}(\text{Div}, \Omega), \\ D(A_1) &= D(\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top) = \mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{RotRot}^\top, \Omega), & D(A_1^*) &= D(\mathbf{RotRot}_{\mathbb{S}, \Gamma_n}^\top) = \mathbf{H}_{\mathbb{S}, \Gamma_n}(\text{RotRot}^\top, \Omega), \\ D(A_2) &= D(\text{Div}_{\mathbb{S}, \Gamma_t}) = \mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{Div}, \Omega), & D(A_2^*) &= D(\mathbf{symGrad}_{\Gamma_n}) = \mathbf{H}_{\Gamma_n}(\text{symGrad}, \Omega). \end{aligned}$$

We shall introduce the latter Sobolev spaces in the next section.

## 2.3 | Sobolev spaces

Let

$$\begin{aligned} \mathbf{H}(\text{symGrad}, \Omega) &:= \{v \in L^2(\Omega) : \text{symGrad} v \in L^2(\Omega)\}, \\ \mathbf{H}_{\mathbb{S}}(\text{RotRot}^\top, \Omega) &:= \{S \in L^2_{\mathbb{S}}(\Omega) : \text{RotRot}^\top S \in L^2(\Omega)\}, \\ \mathbf{H}_{\mathbb{S}}(\text{Div}, \Omega) &:= \{T \in L^2_{\mathbb{S}}(\Omega) : \text{Div} T \in L^2(\Omega)\}. \end{aligned}$$

Note that  $M \in \mathbf{H}_{\mathbb{S}}(\text{RotRot}^\top, \Omega)$  implies  $\text{RotRot}^\top M \in L^2_{\mathbb{S}}(\Omega)$  and that we have by Korn's inequality the regularity

$$\mathbf{H}(\text{symGrad}, \Omega) = \mathbf{H}^1(\Omega)$$

with equivalent norms. Moreover, we define boundary conditions in the *strong sense* as closures of respective test fields, that is,

$$\begin{aligned} \mathbf{H}_{\Gamma_t}(\text{symGrad}, \Omega) &:= \overline{\mathbf{H}(\text{symGrad}, \Omega)}_{\mathbf{C}_{\Gamma_t}^\infty(\Omega)}, \\ \mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{RotRot}^\top, \Omega) &:= \overline{\mathbf{H}_{\mathbb{S}}(\text{RotRot}^\top, \Omega)}_{\mathbf{C}_{\mathbb{S}, \Gamma_t}^\infty(\Omega)}, \\ \mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{Div}, \Omega) &:= \overline{\mathbf{H}_{\mathbb{S}}(\text{Div}, \Omega)}_{\mathbf{C}_{\mathbb{S}, \Gamma_t}^\infty(\Omega)}, \end{aligned}$$

and we have  $\mathbf{H}_{\emptyset}(\text{symGrad}, \Omega) = \mathbf{H}(\text{symGrad}, \Omega) = \mathbf{H}^1(\Omega)$ ,  $\mathbf{H}_{\mathbb{S}, \emptyset}(\text{RotRot}^\top, \Omega) = \mathbf{H}_{\mathbb{S}}(\text{RotRot}^\top, \Omega)$ , and  $\mathbf{H}_{\mathbb{S}, \emptyset}(\text{Div}, \Omega) = \mathbf{H}_{\mathbb{S}}(\text{Div}, \Omega)$ , which are density results and incorporated into the notation by purpose. Spaces with vanishing  $\text{RotRot}^\top$  and  $\text{Div}$  are denoted by  $\mathbf{H}_{\mathbb{S}, \Gamma_t, 0}(\text{RotRot}^\top, \Omega)$  and  $\mathbf{H}_{\mathbb{S}, \Gamma_t, 0}(\text{Div}, \Omega)$ , respectively. Note that, again by Korn's inequality, we have

$$\mathbf{H}_{\Gamma_t}(\text{symGrad}, \Omega) = \mathbf{H}_{\Gamma_t}^1(\Omega).$$

We need also the Sobolev spaces with boundary conditions defined in the *weak sense*, that is,

$$\begin{aligned} \mathbf{H}_{\Gamma_t}(\text{symGrad}, \Omega) &:= \left\{ v \in \mathbf{H}(\text{symGrad}, \Omega) : \langle \text{symGrad} v, \Phi \rangle_{L^2(\Omega)} = -\langle v, \text{Div} \Phi \rangle_{L^2(\Omega)} \forall \Phi \in \mathbf{C}_{\mathbb{S}, \Gamma_n}^\infty(\Omega) \right\}, \\ \mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{RotRot}^\top, \Omega) &:= \left\{ S \in \mathbf{H}_{\mathbb{S}}(\text{RotRot}^\top, \Omega) : \langle \text{RotRot}^\top S, \Psi \rangle_{L^2(\Omega)} = \langle S, \text{RotRot}^\top \Psi \rangle_{L^2(\Omega)} \forall \Psi \in \mathbf{C}_{\mathbb{S}, \Gamma_n}^\infty(\Omega) \right\}, \\ \mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{Div}, \Omega) &:= \left\{ T \in \mathbf{H}_{\mathbb{S}}(\text{Div}, \Omega) : \langle \text{Div} T, \phi \rangle_{L^2(\Omega)} = -\langle T, \text{symGrad} \phi \rangle_{L^2(\Omega)} \forall \phi \in \mathbf{C}_{\Gamma_n}^\infty(\Omega) \right\}. \end{aligned}$$

Note that “*strong*  $\subset$  *weak*” holds; that is,  $\mathbf{H}_{\dots}(\dots, \Omega) \subset \mathbf{H}_{\dots}(\dots, \Omega)$ , for example,

$$\mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{RotRot}^\top, \Omega) \subset \mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{RotRot}^\top, \Omega), \quad \mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{Div}, \Omega) \subset \mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{Div}, \Omega),$$

and that the complex properties hold in both the strong case and the weak case, for example,

$$\text{symGrad } H_{\Gamma_t}(\Omega) \subset H_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^\top, \Omega), \quad \text{RotRot}^\top H_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) \subset H_{\mathbb{S},\Gamma_t,0}(\text{Div}, \Omega),$$

which follows immediately by the definitions. In Remark 2.4, we comment on the question whether “strong = weak” holds in general.

### 2.4 | Higher order Sobolev spaces

For  $k \in \mathbb{N}_0$ , we define higher order Sobolev spaces by

$$\begin{aligned} H_{\mathbb{S}}^k(\Omega) &:= H^k(\Omega) \cap L_{\mathbb{S}}^2(\Omega), \\ H_{\mathbb{S},\Gamma_t}^k(\Omega) &:= \overline{C_{\mathbb{S},\Gamma_t}^\infty(\Omega)}^{H^k(\Omega)} = H_{\Gamma_t}^k(\Omega) \cap L_{\mathbb{S}}^2(\Omega), \\ H^k(\text{symGrad}, \Omega) &:= \left\{ v \in H^k(\Omega) : \text{symGrad } v \in H^k(\Omega) \right\}, \\ H_{\Gamma_t}^k(\text{symGrad}, \Omega) &:= \left\{ v \in H_{\Gamma_t}^k(\Omega) \cap H_{\Gamma_t}(\text{symGrad}, \Omega) : \text{symGrad } v \in H_{\Gamma_t}^k(\Omega) \right\}, \\ H_{\mathbb{S}}^k(\text{RotRot}^\top, \Omega) &:= \left\{ S \in H_{\mathbb{S}}^k(\Omega) : \text{RotRot}^\top S \in H^k(\Omega) \right\}, \\ H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) &:= \left\{ S \in H_{\mathbb{S},\Gamma_t}^k(\Omega) \cap H_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) : \text{RotRot}^\top S \in H_{\Gamma_t}^k(\Omega) \right\}, \\ H_{\mathbb{S}}^k(\text{Div}, \Omega) &:= \left\{ T \in H_{\mathbb{S}}^k(\Omega) : \text{Div } T \in H^k(\Omega) \right\}, \\ H_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) &:= \left\{ T \in H_{\mathbb{S},\Gamma_t}^k(\Omega) \cap H_{\mathbb{S},\Gamma_t}(\text{Div}, \Omega) : \text{Div } T \in H_{\Gamma_t}^k(\Omega) \right\}. \end{aligned}$$

We see  $H_{\mathbb{S},\emptyset}^k(\text{RotRot}^\top, \Omega) = H_{\mathbb{S}}^k(\text{RotRot}^\top, \Omega)$  and  $H_{\mathbb{S},\emptyset}^0(\text{RotRot}^\top, \Omega) = H_{\mathbb{S}}(\text{RotRot}^\top, \Omega)$  as well as  $H_{\mathbb{S},\Gamma_t}^0(\text{RotRot}^\top, \Omega) = H_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega)$ . Note that for  $\Gamma_t \neq \emptyset$  it holds

$$H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) = \left\{ S \in H_{\mathbb{S},\Gamma_t}^k(\Omega) : \text{RotRot}^\top S \in H_{\Gamma_t}^k(\Omega) \right\}, \quad k \geq 2, \tag{2}$$

but for  $\Gamma_t \neq \emptyset$  and  $k = 0$  and  $k = 1$  (as  $H_{\mathbb{S},\Gamma_t}^0(\Omega) = L_{\mathbb{S}}^2(\Omega)$ )

$$\begin{aligned} H_{\mathbb{S},\Gamma_t}^0(\text{RotRot}^\top, \Omega) &= H_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) \\ &\subsetneq \left\{ S \in H_{\mathbb{S},\Gamma_t}^0(\Omega) : \text{RotRot}^\top S \in H_{\Gamma_t}^0(\Omega) \right\} = H_{\mathbb{S}}(\text{RotRot}^\top, \Omega), \\ H_{\mathbb{S},\Gamma_t}^1(\text{RotRot}^\top, \Omega) &\subsetneq \left\{ S \in H_{\mathbb{S},\Gamma_t}^1(\Omega) : \text{RotRot}^\top S \in H_{\Gamma_t}^1(\Omega) \right\}, \end{aligned}$$

respectively. As before, we introduce the kernels

$$\begin{aligned} H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega) &:= H_{\Gamma_t}^k(\Omega) \cap H_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^\top, \Omega) = H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \cap H_{\mathbb{S},0}(\text{RotRot}^\top, \Omega) \\ &= \left\{ S \in H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) : \text{RotRot}^\top S = 0 \right\}. \end{aligned}$$

The corresponding remarks and definitions extend to the  $H_{\Gamma_t}^k(\text{symGrad}, \Omega)$ -spaces and  $H_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega)$ -spaces as well. In particular, we have for  $\Gamma_t \neq \emptyset$  and  $k \geq 1$

$$\begin{aligned} H_{\Gamma_t}^k(\text{symGrad}, \Omega) &= \left\{ v \in H_{\Gamma_t}^k(\Omega) : \text{symGrad } v \in H_{\Gamma_t}^k(\Omega) \right\}, \\ H_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) &= \left\{ T \in H_{\mathbb{S},\Gamma_t}^k(\Omega) : \text{Div } T \in H_{\Gamma_t}^k(\Omega) \right\}, \end{aligned} \tag{3}$$

and

$$\begin{aligned} H_{\Gamma_t}^0(\text{symGrad}, \Omega) &= H_{\Gamma_t}(\text{symGrad}, \Omega) \subsetneq \left\{ v \in H_{\Gamma_t}^0(\Omega) : \text{symGrad} v \in H_{\Gamma_t}^0(\Omega) \right\} = H(\text{symGrad}, \Omega), \\ H_{\mathbb{S}, \Gamma_t}^0(\text{Div}, \Omega) &= H_{\mathbb{S}, \Gamma_t}(\text{Div}, \Omega) \subsetneq \left\{ T \in H_{\mathbb{S}, \Gamma_t}^0(\Omega) : \text{Div} T \in H_{\Gamma_t}^0(\Omega) \right\} = H_{\mathbb{S}}(\text{Div}, \Omega), \end{aligned}$$

as well as

$$\begin{aligned} H_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega) &= H_{\Gamma_t}^k(\Omega) \cap H_{\mathbb{S}, \Gamma_t, 0}(\text{Div}, \Omega) = H_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega) \cap H_{\mathbb{S}, 0}(\text{Div}, \Omega) \\ &= \left\{ T \in H_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega) : \text{Div} T = 0 \right\}. \end{aligned}$$

Analogously, we define the Sobolev spaces  $\mathbf{H}_{\Gamma_t}^k(\text{symGrad}, \Omega)$ ,  $\mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega)$ ,  $\mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega)$ , and  $\mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega)$ ,  $\mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega)$  using the respective Sobolev spaces with weak boundary conditions. Note that again “*strong*  $\subset$  *weak*” holds; that is,  $H_{\mathbb{S}, \Gamma_t}(\dots, \Omega) \subset \mathbf{H}_{\mathbb{S}, \Gamma_t}(\dots, \Omega)$ , for example,

$$H_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega) \subset \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega), \quad H_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega) \subset \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega),$$

and that the complex properties hold in both the strong case and the weak case, for example,

$$\text{symGrad} H_{\Gamma_t}^{k+1}(\Omega) \subset H_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega), \quad \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega) \subset H_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega).$$

Moreover, the corresponding results for (2) and (3) hold also for the weak spaces.

In the forthcoming sections, we shall also investigate whether indeed “*strong* = *weak*” holds. We start with a simple implication from Lemma 2.1.

**Corollary 2.2.**  $H_{\mathbb{S}, \Gamma_t}^k(\Omega) = \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\Omega)$ ; that is, weak and strong boundary conditions coincide for the standard Sobolev spaces of symmetric tensor fields with mixed boundary conditions.

Lemma 2.1, Corollary 2.2, Equations (2) and (3), and Korn’s inequality show the following.

**Lemma 2.3** (higher order weak and strong partial boundary conditions coincide).

- (i) For  $k \geq 0$ , it holds  $H_{\Gamma_t}^k(\text{symGrad}, \Omega) = H_{\Gamma_t}^{k+1}(\Omega) = \mathbf{H}_{\Gamma_t}^{k+1}(\Omega)$ .
- (ii) For  $k \geq 1$ , it holds

$$\begin{aligned} H_{\Gamma_t}^k(\text{symGrad}, \Omega) &= \left\{ v \in H_{\Gamma_t}^k(\Omega) : \text{symGrad} v \in H_{\Gamma_t}^k(\Omega) \right\} = H_{\Gamma_t}^k(\text{symGrad}, \Omega) = H_{\Gamma_t}^{k+1}(\Omega), \\ H_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega) &= \left\{ T \in H_{\mathbb{S}, \Gamma_t}^k(\Omega) : \text{Div} T \in H_{\Gamma_t}^k(\Omega) \right\} = H_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega). \end{aligned}$$

- (iii) For  $k \geq 2$ , it holds

$$H_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega) = \left\{ S \in H_{\mathbb{S}, \Gamma_t}^k(\Omega) : \text{RotRot}^\top S \in H_{\Gamma_t}^k(\Omega) \right\} = \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega).$$

**Remark 2.4** (weak and strong partial boundary conditions coincide). In references,<sup>10,11</sup> we could prove the corresponding results “*strong* = *weak*” for the whole elasticity complex but only with empty or full boundary conditions ( $\Gamma_t = \emptyset$  or  $\Gamma_t = \Gamma$ ). Therefore, in these special cases, the adjoints are well defined on the spaces with strong boundary conditions as well.

Lemma 2.3 shows that for higher values of  $k$  indeed “*strong* = *weak*” holds. Thus, to show “*strong* = *weak*” in general, we only have to prove that equality holds in the remaining cases  $k = 0$  and  $k = 1$ ; that is, we only have to show

$$\begin{aligned} H_{\Gamma_t}(\text{symGrad}, \Omega) &\subset H_{\Gamma_t}(\text{symGrad}, \Omega), & H_{\mathbb{S}, \Gamma_t}(\text{RotRot}^\top, \Omega) &\subset H_{\mathbb{S}, \Gamma_t}(\text{RotRot}^\top, \Omega), \\ H_{\mathbb{S}, \Gamma_t}(\text{Div}, \Omega) &\subset H_{\mathbb{S}, \Gamma_t}(\text{Div}, \Omega), & H_{\mathbb{S}, \Gamma_t}^1(\text{RotRot}^\top, \Omega) &\subset H_{\mathbb{S}, \Gamma_t}^1(\text{RotRot}^\top, \Omega). \end{aligned}$$

The most delicate situation appears due to the second-order nature of  $\text{RotRot}_{\mathbb{S}}^{\top}$ . In Corollary 3.11, we shall show using regular decompositions that these results (weak and strong boundary conditions coincide for the elasticity complex for all  $k \geq 0$ ) indeed hold true.

### 2.5 | More Sobolev spaces

For  $k \in \mathbb{N}$ , we introduce also slightly less regular higher order Sobolev spaces by

$$\begin{aligned} \mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^{\top}, \Omega) &:= \left\{ S \in \mathbf{H}_{\Gamma_t}^k(\Omega) \cap \mathbf{H}_{\mathbb{S},\Gamma_t}(\text{RotRot}^{\top}, \Omega) : \text{RotRot}^{\top} S \in \mathbf{H}_{\Gamma_t}^{k-1}(\Omega) \right\}, \\ \mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^{\top}, \Omega) &:= \left\{ S \in \mathbf{H}_{\Gamma_t}^k(\Omega) \cap \mathbf{H}_{\mathbb{S},\Gamma_t}(\text{RotRot}^{\top}, \Omega) : \text{RotRot}^{\top} S \in \mathbf{H}_{\Gamma_t}^{k-1}(\Omega) \right\}, \end{aligned}$$

and we extend all conventions of our notations. For the kernels, we have

$$\mathbf{H}_{\mathbb{S},\Gamma_t,0}^{k,k-1}(\text{RotRot}^{\top}, \Omega) = \mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^{\top}, \Omega), \quad \mathbf{H}_{\mathbb{S},\Gamma_t,0}^{k,k-1}(\text{RotRot}^{\top}, \Omega) = \mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^{\top}, \Omega).$$

Note that, as before, the intersection with  $\mathbf{H}_{\mathbb{S},\Gamma_t}(\text{RotRot}^{\top}, \Omega)$  and  $\mathbf{H}_{\mathbb{S},\Gamma_t}(\text{RotRot}^{\top}, \Omega)$  is only needed if  $k = 1$ . Again, we have “*strong*  $\subset$  *weak*,” that is,  $\mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^{\top}, \Omega) \subset \mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^{\top}, \Omega)$ , and in both cases (weak and strong), the complex properties hold, that is,

$$\text{symGrad } \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \subset \mathbf{H}_{\mathbb{S},\Gamma_t,0}^{k,k-1}(\text{RotRot}^{\top}, \Omega), \quad \text{RotRot}^{\top} \mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^{\top}, \Omega) \subset \mathbf{H}_{\mathbb{S},\Gamma_t,0}^{k-1}(\text{Div}, \Omega).$$

Similar to Lemma 2.3, we have the following.

**Lemma 2.5** (higher order weak and strong partial boundary conditions coincide). *For  $k \geq 2$ ,*

$$\mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^{\top}, \Omega) = \left\{ S \in \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\Omega) : \text{RotRot}^{\top} S \in \mathbf{H}_{\Gamma_t}^{k-1}(\Omega) \right\} = \mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^{\top}, \Omega).$$

### 2.6 | Some elasticity complexes

By definition, we have densely defined and closed (unbounded) linear operators defining three dual pairs

$$\begin{aligned} (\text{symGrad}_{\Gamma_t}, \text{symGrad}_{\Gamma_t}^*) &= (\text{symGrad}_{\Gamma_t}, -\mathbf{Div}_{\mathbb{S},\Gamma_n}), \\ (\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top}, (\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top})^*) &= (\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top}, \text{RotRot}_{\mathbb{S},\Gamma_n}^{\top}), \\ (\text{Div}_{\mathbb{S},\Gamma_t}, \text{Div}_{\mathbb{S},\Gamma_t}^*) &= (\text{Div}_{\mathbb{S},\Gamma_t}, -\mathbf{symGrad}_{\Gamma_n}). \end{aligned}$$

Remarks 2.5 and 2.6<sup>1,2</sup> show the complex properties

$$\begin{aligned} \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top} \text{symGrad}_{\Gamma_t} &\subset 0, & \text{Div}_{\mathbb{S},\Gamma_t} \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top} &\subset 0, \\ -\mathbf{Div}_{\mathbb{S},\Gamma_n} \text{RotRot}_{\mathbb{S},\Gamma_n}^{\top} &\subset 0, & -\text{RotRot}_{\mathbb{S},\Gamma_n}^{\top} \mathbf{symGrad}_{\Gamma_n} &\subset 0. \end{aligned}$$

Hence, we get the primal and dual elasticity Hilbert complex

$$\begin{array}{ccccccc} \dots & \dots & \text{symGrad}_{\Gamma_t} & & \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top} & & \text{Div}_{\mathbb{S},\Gamma_t} & \dots \\ \dots & \rightleftarrows & L^2(\Omega) & \rightleftarrows & L^2_{\mathbb{S}}(\Omega) & \rightleftarrows & L^2(\Omega) & \rightleftarrows \dots \\ \dots & & -\mathbf{Div}_{\mathbb{S},\Gamma_n} & & \text{RotRot}_{\mathbb{S},\Gamma_n}^{\top} & & -\mathbf{symGrad}_{\Gamma_n} & \dots \end{array} \tag{4}$$

with the complex properties

$$\begin{aligned} R(\text{symGrad}_{\Gamma_t}) &\subset N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top), & R(\mathbf{RotRot}_{\mathbb{S},\Gamma_n}^\top) &\subset N(\mathbf{Div}_{\mathbb{S},\Gamma_n}), \\ R(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top) &\subset N(\mathbf{Div}_{\mathbb{S},\Gamma_t}), & R(\mathbf{symGrad}_{\Gamma_n}) &\subset N(\mathbf{RotRot}_{\mathbb{S},\Gamma_n}^\top). \end{aligned}$$

The long primal and dual elasticity Hilbert complex (cf. eq. 12<sup>1,2</sup>) reads

$$\mathbb{RM}_{\Gamma_t} \begin{array}{c} \xleftarrow{\text{'}\mathbb{RM}_{\Gamma_t}} \\ \xrightarrow{\text{'}\mathbb{RM}_{\Gamma_t}} \end{array} L^2(\Omega) \begin{array}{c} \xrightarrow{\text{symGrad}_{\Gamma_t}} \\ \xleftarrow{-\mathbf{Div}_{\mathbb{S},\Gamma_n}} \end{array} L^2_{\mathbb{S}}(\Omega) \begin{array}{c} \xrightarrow{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top} \\ \xleftarrow{\mathbf{RotRot}_{\mathbb{S},\Gamma_n}^\top} \end{array} L^2_{\mathbb{S}}(\Omega) \begin{array}{c} \xrightarrow{\mathbf{Div}_{\mathbb{S},\Gamma_t}} \\ \xleftarrow{-\mathbf{symGrad}_{\Gamma_n}} \end{array} L^2(\Omega) \begin{array}{c} \xrightarrow{\text{'}\mathbb{RM}_{\Gamma_n}} \\ \xleftarrow{\text{'}\mathbb{RM}_{\Gamma_n}} \end{array} \mathbb{RM}_{\Gamma_n} \quad (5)$$

with the additional complex properties

$$\begin{aligned} R(\iota_{\mathbb{RM}_{\Gamma_t}}) &= N(\text{symGrad}_{\Gamma_t}) = \mathbb{RM}_{\Gamma_t}, & \overline{R(\mathbf{Div}_{\mathbb{S},\Gamma_n})} &= \mathbb{RM}_{\Gamma_t}^{\perp L^2(\Omega)}, \\ \overline{R(\mathbf{Div}_{\mathbb{S},\Gamma_t})} &= \mathbb{RM}_{\Gamma_n}^{\perp L^2(\Omega)}, & R(\iota_{\mathbb{RM}_{\Gamma_n}}) &= N(\mathbf{symGrad}_{\Gamma_n}) = \mathbb{RM}_{\Gamma_n}, \end{aligned}$$

where

$$\mathbb{RM}_{\Sigma} = \begin{cases} \{0\} & \text{if } \Sigma \neq \emptyset, \\ \mathbb{RM} & \text{if } \Sigma = \emptyset, \end{cases} \quad \text{with} \quad \mathbb{RM} := \{x \mapsto Qx + q : Q \in \mathbb{R}^{3 \times 3} \text{ skew}, q \in \mathbb{R}^3\}$$

denoting the global rigid motions in  $\Omega$ . Note that  $\dim \mathbb{RM} = 6$ .

More generally, in addition to (5), we shall discuss for  $k \in \mathbb{N}_0$  the higher Sobolev order (long primal and formally dual) elasticity Hilbert complexes (omitting  $\Omega$  in the notation)

$$\begin{array}{ccccccccccc} \mathbb{RM}_{\Gamma_t} & \xrightarrow{\text{'}\mathbb{RM}_{\Gamma_t}} & H_{\Gamma_t}^k & \xrightarrow{\text{symGrad}_{\Gamma_t}^k} & H_{\mathbb{S},\Gamma_t}^k & \xrightarrow{\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}} & H_{\mathbb{S},\Gamma_t}^k & \xrightarrow{\mathbf{Div}_{\mathbb{S},\Gamma_t}^k} & H_{\Gamma_t}^k & \xrightarrow{\text{'}\mathbb{RM}_{\Gamma_n}} & \mathbb{RM}_{\Gamma_n} \\ \mathbb{RM}_{\Gamma_t} & \xleftarrow{\text{'}\mathbb{RM}_{\Gamma_t}} & H_{\Gamma_n}^k & \xleftarrow{-\mathbf{Div}_{\mathbb{S},\Gamma_n}^k} & H_{\mathbb{S},\Gamma_n}^k & \xleftarrow{\mathbf{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k}} & H_{\mathbb{S},\Gamma_n}^k & \xleftarrow{-\mathbf{symGrad}_{\Gamma_n}^k} & H_{\Gamma_n}^k & \xleftarrow{\text{'}\mathbb{RM}_{\Gamma_n}} & \mathbb{RM}_{\Gamma_n} \end{array}$$

with associated domain complexes

$$\begin{array}{ccccccccccc} \mathbb{RM}_{\Gamma_t} & \xrightarrow{\text{'}\mathbb{RM}_{\Gamma_t}} & H_{\Gamma_t}^k(\text{symGrad}) & \xrightarrow{\text{symGrad}_{\Gamma_t}^k} & H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top) & \xrightarrow{\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}} & H_{\mathbb{S},\Gamma_t}^k(\text{Div}) & \xrightarrow{\mathbf{Div}_{\mathbb{S},\Gamma_t}^k} & H_{\Gamma_t}^k & \xrightarrow{\text{'}\mathbb{RM}_{\Gamma_n}} & \mathbb{RM}_{\Gamma_n} \\ \mathbb{RM}_{\Gamma_t} & \xleftarrow{\text{'}\mathbb{RM}_{\Gamma_t}} & H_{\Gamma_n}^k & \xleftarrow{-\mathbf{Div}_{\mathbb{S},\Gamma_n}^k} & H_{\mathbb{S},\Gamma_n}^k(\text{Div}) & \xleftarrow{\mathbf{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k}} & H_{\mathbb{S},\Gamma_n}^k(\text{RotRot}^\top) & \xleftarrow{-\mathbf{symGrad}_{\Gamma_n}^k} & H_{\Gamma_n}^k(\text{symGrad}) & \xleftarrow{\text{'}\mathbb{RM}_{\Gamma_n}} & \mathbb{RM}_{\Gamma_n} \end{array}$$

Additionally, for  $k \geq 1$ , we will also discuss the following variants of the elasticity complexes:

$$\begin{array}{ccccccccccc} \mathbb{RM}_{\Gamma_t} & \xrightarrow{\text{'}\mathbb{RM}_{\Gamma_t}} & H_{\Gamma_t}^k & \xrightarrow{\text{symGrad}_{\Gamma_t}^k} & H_{\mathbb{S},\Gamma_t}^k & \xrightarrow{\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k,k-1}} & H_{\mathbb{S},\Gamma_t}^{k-1} & \xrightarrow{\mathbf{Div}_{\mathbb{S},\Gamma_t}^{k-1}} & H_{\Gamma_t}^{k-1} & \xrightarrow{\text{'}\mathbb{RM}_{\Gamma_n}} & \mathbb{RM}_{\Gamma_n} \\ \mathbb{RM}_{\Gamma_t} & \xleftarrow{\text{'}\mathbb{RM}_{\Gamma_t}} & H_{\Gamma_n}^{k-1} & \xleftarrow{-\mathbf{Div}_{\mathbb{S},\Gamma_n}^{k-1}} & H_{\mathbb{S},\Gamma_n}^{k-1} & \xleftarrow{\mathbf{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k,k-1}} & H_{\mathbb{S},\Gamma_n}^k & \xleftarrow{-\mathbf{symGrad}_{\Gamma_n}^k} & H_{\Gamma_n}^k & \xleftarrow{\text{'}\mathbb{RM}_{\Gamma_n}} & \mathbb{RM}_{\Gamma_n} \end{array}$$



with associated domain complexes

$$\begin{array}{ccccccccccc}
 \mathbb{R}M_{\Gamma_t} & \xrightarrow{\iota\mathbb{R}M_{\Gamma_t}} & H_{\Gamma_t}^k(\text{symGrad}) & \xrightarrow{\text{symGrad}_{\Gamma_t}^k} & H_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top) & \xrightarrow{\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k,k-1}} & H_{\mathbb{S},\Gamma_t}^{k-1}(\text{Div}) & \xrightarrow{\text{Div}_{\mathbb{S},\Gamma_t}^{k-1}} & H_{\Gamma_t}^{k-1} & \xrightarrow{\pi\mathbb{R}M_{\Gamma_n}} & \mathbb{R}M_{\Gamma_n}, \\
 \mathbb{R}M_{\Gamma_t} & \xleftarrow{\pi\mathbb{R}M_{\Gamma_t}} & H_{\Gamma_t}^{k-1} & \xleftarrow{-\text{Div}_{\mathbb{S},\Gamma_n}^{k-1}} & H_{\mathbb{S},\Gamma_n}^{k-1}(\text{Div}) & \xleftarrow{\text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k,k-1}} & H_{\mathbb{S},\Gamma_n}^{k,k-1}(\text{RotRot}^\top) & \xleftarrow{-\text{symGrad}_{\Gamma_n}^k} & H_{\Gamma_n}^k(\text{symGrad}) & \xleftarrow{\iota\mathbb{R}M_{\Gamma_n}} & \mathbb{R}M_{\Gamma_n}.
 \end{array}$$

Here, we have introduced the densely defined and closed linear operators

$$\begin{aligned}
 \text{symGrad}_{\Gamma_t}^k &: D(\text{symGrad}_{\Gamma_t}^k) \subset H_{\Gamma_t}^k(\Omega) \rightarrow H_{\mathbb{S},\Gamma_t}^k(\Omega); \quad v \mapsto \text{symGrad } v, \\
 \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k} &: D(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}) \subset H_{\mathbb{S},\Gamma_t}^k(\Omega) \rightarrow H_{\mathbb{S},\Gamma_t}^k(\Omega); \quad S \mapsto \text{RotRot}^\top S, \\
 \text{Div}_{\mathbb{S},\Gamma_t}^k &: D(\text{Div}_{\mathbb{S},\Gamma_t}^k) \subset H_{\mathbb{S},\Gamma_t}^k(\Omega) \rightarrow H_{\Gamma_t}^k(\Omega); \quad T \mapsto \text{Div } T, \\
 -\text{Div}_{\mathbb{S},\Gamma_n}^k &: D(\text{Div}_{\mathbb{S},\Gamma_n}^k) \subset H_{\mathbb{S},\Gamma_n}^k(\Omega) \rightarrow H^k\Gamma_n(\Omega); \quad S \mapsto -\text{Div } S, \\
 \text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k} &: D(\text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k}) \subset H_{\mathbb{S},\Gamma_n}^k(\Omega) \rightarrow H_{\mathbb{S},\Gamma_n}^k(\Omega); \quad T \mapsto \text{RotRot}^\top T, \\
 -\text{symGrad}_{\Gamma_n}^k &: D(\text{symGrad}_{\Gamma_n}^k) \subset H_{\Gamma_n}^k(\Omega) \rightarrow H_{\mathbb{S},\Gamma_n}^k(\Omega); \quad v \mapsto -\text{symGrad } v,
 \end{aligned}$$

with domains of definition

$$\begin{aligned}
 D(\text{symGrad}_{\Gamma_t}^k) &= H_{\Gamma_t}^k(\text{symGrad}, \Omega), & D(\text{Div}_{\mathbb{S},\Gamma_n}^k) &= H_{\mathbb{S},\Gamma_n}^k(\text{Div}, \Omega), \\
 D(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}) &= H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega), & D(\text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k}) &= H_{\mathbb{S},\Gamma_n}^k(\text{RotRot}^\top, \Omega), \\
 D(\text{Div}_{\mathbb{S},\Gamma_t}^k) &= H_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega), & D(\text{symGrad}_{\Gamma_n}^k) &= H_{\Gamma_n}^k(\text{symGrad}, \Omega).
 \end{aligned}$$

Moreover,

$$\begin{aligned}
 \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k,k-1} &: D(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k,k-1}) \subset H_{\mathbb{S},\Gamma_t}^k(\Omega) \rightarrow H_{\mathbb{S},\Gamma_t}^{k-1}(\Omega); & S &\mapsto \text{RotRot}^\top S, \\
 \text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k,k-1} &: D(\text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k,k-1}) \subset H_{\mathbb{S},\Gamma_n}^k(\Omega) \rightarrow H_{\mathbb{S},\Gamma_n}^{k-1}(\Omega); & T &\mapsto \text{RotRot}^\top T,
 \end{aligned}$$

with domains of definition

$$D(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k,k-1}) = H_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega), \quad D(\text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k,k-1}) = H_{\mathbb{S},\Gamma_n}^{k,k-1}(\text{RotRot}^\top, \Omega).$$

### 2.7 | Dirichlet/Neumann fields

We also introduce the cohomology space of elastic Dirichlet/Neumann tensor fields (generalized harmonic tensors)

$$\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) := N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top) \cap N(\text{Div}_{\mathbb{S},\Gamma_n}\varepsilon) = H_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^\top, \Omega) \cap \varepsilon^{-1}H_{\mathbb{S},\Gamma_n,0}(\text{Div}, \Omega).$$

Here,  $\varepsilon : L_{\mathbb{S}}^2(\Omega) \rightarrow L_{\mathbb{S}}^2(\Omega)$  is a symmetric and positive topological isomorphism (symmetric and positive bijective bounded linear operator), which introduces a new inner product

$$\langle \cdot, \cdot \rangle_{L_{\mathbb{S},\varepsilon}^2(\Omega)} := \langle \varepsilon \cdot, \cdot \rangle_{L_{\mathbb{S}}^2(\Omega)},$$

where  $L_{\mathbb{S},\varepsilon}^2(\Omega) := L_{\mathbb{S}}^2(\Omega)$  (as linear space) equipped with the inner product  $\langle \cdot, \cdot \rangle_{L_{\mathbb{S},\varepsilon}^2(\Omega)}$ . Such *weights*  $\varepsilon$  shall be called *admissible* and a typical example is given by a symmetric,  $L^\infty$ -bounded, and uniformly positive definite tensor field  $\varepsilon : \Omega \rightarrow \mathbb{R}^{(3 \times 3) \times (3 \times 3)}$ .

### 3 | ELASTICITY COMPLEXES II

#### 3.1 | Regular potentials and decompositions I

##### 3.1.1 | Extendable domains

**Theorem 3.1** (regular potential operators for extendable domains). *Let  $(\Omega, \Gamma_t)$  be an extendable bounded strong Lipschitz pair and let  $k \geq 0$ . Then there exist bounded linear regular potential operators*

$$\begin{aligned} \mathcal{P}_{\text{symGrad}, \Gamma_t}^k &: \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \cap \mathbf{H}^{k+1}(\mathbb{R}^3), \\ \mathcal{P}_{\text{RotRot}^\top, \Gamma_t}^k &: \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega) \rightarrow \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) \cap \mathbf{H}^{k+2}(\mathbb{R}^3), \\ \mathcal{P}_{\text{Div}, \Gamma_t}^k &: \mathbf{H}_{\Gamma_t}^k(\Omega) \cap (\mathbb{R}\mathbb{M}_{\Gamma_n})^{\perp L^2(\Omega)} \rightarrow \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega) \cap \mathbf{H}^{k+1}(\mathbb{R}^3). \end{aligned}$$

In particular,  $\mathcal{P}^{\dots}$  are right inverses for  $\text{symGrad}$ ,  $\text{RotRot}^\top$ , and  $\text{Div}$ , respectively, that is,

$$\begin{aligned} \text{symGrad} \mathcal{P}_{\text{symGrad}, \Gamma_t}^k &= \text{id}_{\mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega)}, \\ \text{RotRot}^\top \mathcal{P}_{\text{RotRot}^\top, \Gamma_t}^k &= \text{id}_{\mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega)}, \\ \text{Div} \mathcal{P}_{\text{Div}, \Gamma_t}^k &= \text{id}_{\mathbf{H}_{\Gamma_t}^k(\Omega) \cap (\mathbb{R}\mathbb{M}_{\Gamma_n})^{\perp L^2(\Omega)}}. \end{aligned}$$

Without loss of generality,  $\mathcal{P}^{\dots}$  map to tensor fields with a fixed compact support in  $\mathbb{R}^3$ .

**Remark 3.2.** Note that  $A_n \mathcal{P}_{A_n} = \text{id}_{R(A_n)}$  is a general property of a (bounded regular) potential operator  $\mathcal{P}_{A_n} : R(A_n) \rightarrow \mathbf{H}_n^+$  with  $\mathbf{H}_n^+ \subset D(A_n)$ .

*Proof of Theorem 3.1.* In theorem 4.2,<sup>10,11</sup> we have shown the stated results for  $\Gamma_t = \Gamma$  and  $\Gamma_t = \emptyset$ , which is also a crucial ingredient of this proof. Note that in these two special cases, always “strong = weak” holds as  $A_n^{**} = \overline{A_n} = A_n$  and that this argument fails in the remaining cases of mixed boundary conditions. Therefore, let  $\emptyset \subsetneq \Gamma_t \subsetneq \Gamma$ . Moreover, recall the notion of an extendable domain from section 3.1.<sup>2</sup> In particular,  $\hat{\Omega}$  and the extended domain  $\tilde{\Omega}$  are topologically trivial.

- Let  $S \in \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega)$ . By definition,  $S$  can be extended through  $\Gamma_t$  by zero to the larger domain  $\tilde{\Omega}$  yielding

$$\tilde{S} \in \mathbf{H}_{\mathbb{S}, \emptyset, 0}^k(\text{RotRot}^\top, \tilde{\Omega}) = \mathbf{H}_{\mathbb{S}, 0}^k(\text{RotRot}^\top, \tilde{\Omega}) = \mathbf{H}_{\mathbb{S}, 0}^k(\text{RotRot}^\top, \tilde{\Omega}).$$

By theorem 4.2,<sup>10,11</sup> there exists  $\tilde{v} \in \mathbf{H}^{k+1}(\mathbb{R}^3)$  such that  $\text{symGrad} \tilde{v} = \tilde{S}$  in  $\tilde{\Omega}$ . Since  $\tilde{S} = 0$  in  $\hat{\Omega}$ ,  $\tilde{v}$  must be a rigid motion  $r \in \mathbb{R}\mathbb{M}$  in  $\hat{\Omega}$ . Far outside of  $\tilde{\Omega}$ , we modify  $r$  by a cut-off function such that the resulting vector field  $\tilde{r}$  is compactly supported and  $\tilde{r}|_{\hat{\Omega}} = r$ . Then  $v := \tilde{v} - \tilde{r} \in \mathbf{H}^{k+1}(\mathbb{R}^3)$  with  $v|_{\hat{\Omega}} = 0$ . Hence,  $v$  belongs to  $\mathbf{H}_{\Gamma_t}^{k+1}(\Omega)$  and depends continuously on  $S$ . Moreover,  $v$  satisfies  $\text{symGrad} v = \text{symGrad} \tilde{v} = \tilde{S}$  in  $\tilde{\Omega}$ , in particular  $\text{symGrad} v = S$  in  $\Omega$ . We put  $\mathcal{P}_{\text{symGrad}, \Gamma_t}^k S := v \in \mathbf{H}_{\Gamma_t}^{k+1}(\Omega)$ .

- Let  $T \in \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega)$ . By definition,  $T$  can be extended through  $\Gamma_t$  by zero to  $\tilde{\Omega}$  giving

$$\tilde{T} \in \mathbf{H}_{\mathbb{S}, \emptyset, 0}^k(\text{Div}, \tilde{\Omega}) = \mathbf{H}_{\mathbb{S}, 0}^k(\text{Div}, \tilde{\Omega}) = \mathbf{H}_{\mathbb{S}, 0}^k(\text{Div}, \tilde{\Omega}).$$

By theorem 4.2,<sup>10,11</sup> there exists  $\tilde{S} \in \mathbf{H}_{\mathbb{S}}^{k+2}(\mathbb{R}^3)$  such that  $\text{RotRot}^\top \tilde{S} = \tilde{T}$  in  $\tilde{\Omega}$ . Since  $\tilde{T} = 0$  in  $\hat{\Omega}$ , i.e.,  $\tilde{S}|_{\hat{\Omega}} \in \mathbf{H}_{\mathbb{S}, 0}^{k+2}(\text{RotRot}^\top, \hat{\Omega})$ , we get again by theorem 4.2<sup>10,11</sup> (or the first part of this proof)  $\tilde{v} \in \mathbf{H}^{k+3}(\mathbb{R}^3)$  such that  $\text{symGrad} \tilde{v} = \tilde{S}$  in  $\hat{\Omega}$ . Then  $S := \tilde{S} - \text{symGrad} \tilde{v}$  belongs to  $\mathbf{H}_{\mathbb{S}}^{k+2}(\mathbb{R}^3)$  and satisfies  $S|_{\hat{\Omega}} = 0$ . Thus,  $S \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega)$  and depends continuously on  $T$ . Furthermore,  $\text{RotRot}^\top S = \text{RotRot}^\top \tilde{S} = \tilde{T}$  in  $\tilde{\Omega}$ , in particular  $\text{RotRot}^\top S = T$  in  $\Omega$ . We set  $\mathcal{P}_{\text{RotRot}^\top, \Gamma_t}^k T := S \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega)$ .

- Let  $v \in H^k_{\Gamma_t}(\Omega)$ . By definition,  $v$  can be extended through  $\Gamma_t$  by zero to  $\tilde{\Omega}$  defining  $\tilde{v} \in H^k(\tilde{\Omega})$ . Theorem 4.2<sup>10,11</sup> yields  $\tilde{T} \in H^{k+1}_{\mathbb{S}}(\mathbb{R}^3)$  such that  $\text{Div} \tilde{T} = \tilde{v}$  in  $\tilde{\Omega}$ . As  $\tilde{v} = 0$  in  $\hat{\Omega}$ , i.e.,  $\tilde{T}|_{\hat{\Omega}} \in H^{k+1}_{\mathbb{S},0}(\text{Div}, \hat{\Omega})$ , we get again by theorem 4.2<sup>10,11</sup> (or the second part of this proof)  $\tilde{S} \in H^{k+3}_{\mathbb{S}}(\mathbb{R}^3)$  such that  $\text{RotRot}^{\top} \tilde{S} = \tilde{T}$  holds in  $\hat{\Omega}$ . Then  $T := \tilde{T} - \text{RotRot}^{\top} \tilde{S}$  belongs to  $H^{k+1}_{\mathbb{S}}(\mathbb{R}^3)$  with  $T|_{\hat{\Omega}} = 0$ . Hence,  $T$  belongs to  $H^{k+1}_{\mathbb{S},\Gamma_t}(\Omega)$  and depends continuously on  $v$ . Furthermore,  $\text{Div} T = \text{Div} \tilde{T} = \tilde{v}$  in  $\tilde{\Omega}$ , in particular  $\text{Div} T = v$  in  $\Omega$ . Finally, we define  $\mathcal{P}^k_{\text{Div}_{\mathbb{S},\Gamma_t}} v := T \in H^{k+1}_{\mathbb{S},\Gamma_t}(\Omega)$ .

The assertion about the compact supports is trivial. □

As a simple consequence of Theorem 3.1, we obtain a few corollaries.

**Corollary 3.3** (regular potentials for extendable domains). *Let  $(\Omega, \Gamma_t)$  be an extendable bounded strong Lipschitz pair and let  $k \geq 0$ . Then the regular potential representations*

$$\begin{aligned} H^k_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^{\top}, \Omega) &= H^k_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^{\top}, \Omega) = \text{symGrad } H^k_{\Gamma_t}(\text{symGrad}, \Omega) = \text{symGrad } H^{k+1}_{\Gamma_t}(\Omega) \\ &= R(\text{symGrad}^k_{\Gamma_t}), \\ H^k_{\mathbb{S},\Gamma_t,0}(\text{Div}, \Omega) &= H^k_{\mathbb{S},\Gamma_t,0}(\text{Div}, \Omega) = \text{RotRot}^{\top} H^k_{\mathbb{S},\Gamma_t}(\text{RotRot}^{\top}, \Omega) = \text{RotRot}^{\top} H^{k+2}_{\mathbb{S},\Gamma_t}(\Omega) \\ &= \text{RotRot}^{\top} H^{k+1,k}_{\mathbb{S},\Gamma_t}(\text{RotRot}^{\top}, \Omega) \\ &= R(\text{RotRot}^{\top,k}_{\mathbb{S},\Gamma_t}) = R(\text{RotRot}^{\top,k+1,k}_{\mathbb{S},\Gamma_t}), \\ H^k_{\Gamma_t}(\Omega) \cap (\mathbb{R}M_{\Gamma_t})^{\perp L^2(\Omega)} &= \text{Div} H^k_{\mathbb{S},\Gamma_t}(\text{Div}, \Omega) = \text{Div} H^{k+1}_{\mathbb{S},\Gamma_t}(\Omega) \\ &= R(\text{Div}^k_{\mathbb{S},\Gamma_t}) \end{aligned}$$

hold, and the potentials can be chosen such that they depend continuously on the data. In particular, the latter spaces are closed subspaces of  $H^k_{\mathbb{S}}(\Omega)$  and  $H^k(\Omega)$ , respectively.

*Proof.* By Theorem 3.1, we have

$$\begin{aligned} H^k_{\mathbb{S},\Gamma_t,0}(\text{Div}, \Omega) &= \text{RotRot}^{\top} \mathcal{P}^k_{\text{RotRot}^{\top}_{\mathbb{S},\Gamma_t}} H^k_{\mathbb{S},\Gamma_t,0}(\text{Div}, \Omega) \subset \text{RotRot}^{\top} H^{k+2}_{\mathbb{S},\Gamma_t}(\Omega) \\ &\subset \text{RotRot}^{\top} H^{k+1,k}_{\mathbb{S},\Gamma_t}(\text{RotRot}^{\top}, \Omega) \subset \text{RotRot}^{\top} H^k_{\mathbb{S},\Gamma_t}(\text{RotRot}^{\top}, \Omega) \\ &\subset H^k_{\mathbb{S},\Gamma_t,0}(\text{Div}, \Omega) \subset H^k_{\mathbb{S},\Gamma_t,0}(\text{Div}, \Omega). \end{aligned}$$

The other identities follow analogously. □

**Corollary 3.4** (regular decompositions for extendable domains). *Let  $(\Omega, \Gamma_t)$  be an extendable bounded strong Lipschitz pair and let  $k \geq 0$ . Then the bounded regular decompositions*

$$\begin{aligned} H^k_{\mathbb{S},\Gamma_t}(\text{RotRot}^{\top}, \Omega) &= H^{k+2}_{\mathbb{S},\Gamma_t}(\Omega) + \text{symGrad } H^{k+1}_{\Gamma_t}(\Omega) = R(\mathcal{P}^k_{\text{RotRot}^{\top}_{\mathbb{S},\Gamma_t}}) \dot{+} H^k_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^{\top}, \Omega) \\ &= R(\mathcal{P}^k_{\text{RotRot}^{\top}_{\mathbb{S},\Gamma_t}}) \dot{+} \text{symGrad } H^{k+1}_{\Gamma_t}(\Omega) \\ &= R(\mathcal{P}^k_{\text{RotRot}^{\top}_{\mathbb{S},\Gamma_t}}) \dot{+} \text{symGrad } R(\mathcal{P}^k_{\text{symGrad},\Gamma_t}), \\ H^k_{\mathbb{S},\Gamma_t}(\text{Div}, \Omega) &= H^{k+1}_{\mathbb{S},\Gamma_t}(\Omega) + \text{RotRot}^{\top} H^{k+2}_{\mathbb{S},\Gamma_t}(\Omega) = R(\mathcal{P}^k_{\text{Div}_{\mathbb{S},\Gamma_t}}) \dot{+} H^k_{\mathbb{S},\Gamma_t,0}(\text{Div}, \Omega) \\ &= R(\mathcal{P}^k_{\text{Div}_{\mathbb{S},\Gamma_t}}) \dot{+} \text{RotRot}^{\top} H^{k+2}_{\mathbb{S},\Gamma_t}(\Omega) \\ &= R(\mathcal{P}^k_{\text{Div}_{\mathbb{S},\Gamma_t}}) \dot{+} \text{RotRot}^{\top} R(\mathcal{P}^k_{\text{RotRot}^{\top}_{\mathbb{S},\Gamma_t}}) \end{aligned}$$

hold with bounded linear regular decomposition operators

$$\begin{aligned} Q_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1} &:= \mathcal{P}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^k \text{RotRot}^\top : \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), \\ Q_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,0} &:= \mathcal{P}_{\text{symGrad},\Gamma_t}^k (1 - Q_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1}) : \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\Gamma_t}^{k+1}(\Omega), \\ Q_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,1} &:= \mathcal{P}_{\text{Div}_{\mathbb{S},\Gamma_t}}^k \text{Div} : \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) \rightarrow \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega), \\ Q_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,0} &:= \mathcal{P}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^k (1 - Q_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,1}) : \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) \rightarrow \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}(\Omega) \end{aligned}$$

satisfying

$$\begin{aligned} Q_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1} + \text{symGrad} Q_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,0} &= \text{id}_{\mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega)}, \\ Q_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,1} + \text{RotRot}^\top Q_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,0} &= \text{id}_{\mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega)}. \end{aligned}$$

*Remark 3.5.* Note that for (bounded linear) potential operators  $\mathcal{P}_{A_{n-1}}$  and  $\mathcal{P}_{A_n}$ , the identity

$$\begin{aligned} Q_{A_n}^1 + A_{n-1} Q_{A_n}^0 &= \text{id}_{D(A_n)} \quad \text{with} \quad Q_{A_n}^1 := \mathcal{P}_{A_n} A_n : D(A_n) \rightarrow \mathbf{H}_n^+, \\ Q_{A_n}^0 &:= \mathcal{P}_{A_{n-1}} (1 - Q_{A_n}^1) : D(A_n) \rightarrow \mathbf{H}_{n-1}^+ \end{aligned}$$

is a general structure of a (bounded) regular decomposition. Moreover,

- (i)  $R(Q_{A_n}^1) = R(\mathcal{P}_{A_n})$  and  $R(Q_{A_n}^0) = R(\mathcal{P}_{A_{n-1}})$ .
- (ii)  $N(A_n)$  is invariant under  $Q_{A_n}^1$ , as  $A_n = A_n Q_{A_n}^1$  holds by the complex property.
- (iii)  $Q_{A_n}^1$  and  $A_{n-1} Q_{A_n}^0 = 1 - Q_{A_n}^1$  are projections.
- (iv) There exists  $c > 0$  such that for all  $x \in D(A_n)$

$$|Q_{A_n}^1 x|_{\mathbf{H}_n^+} \leq c |A_n x|_{\mathbf{H}_{n+1}}.$$

(iv') In particular,  $Q_{A_n}^1|_{N(A_n)} = 0$ .

**Corollary 3.6** (weak and strong partial boundary conditions coincide for extendable domains). *Let  $(\Omega, \Gamma_t)$  be an extendable bounded strong Lipschitz pair and let  $k \geq 0$ . Then weak and strong boundary conditions coincide, that is,*

$$\begin{aligned} \mathbf{H}_{\Gamma_t}^k(\text{symGrad}, \Omega) &= \mathbf{H}_{\Gamma_t}^k(\text{symGrad}, \Omega) = \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) = \mathbf{H}_{\Gamma_t}^{k+1}(\Omega), \\ \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) &= \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega), \\ \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) &= \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega). \end{aligned}$$

*Proof of Corollaries 3.4 and 3.6.* Let us pick  $S \in \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega)$ . By Theorem 3.1, we have  $\text{RotRot}^\top S \in \mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{Div}, \Omega)$  and  $\hat{S} := \mathcal{P}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^k \text{RotRot}^\top S \in \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}$ . Hence, we obtain  $S - \hat{S} \in \mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega)$  and Theorem 3.1 shows  $v := \mathcal{P}_{\text{symGrad},\Gamma_t}^k (S - \hat{S}) \in \mathbf{H}_{\Gamma_t}^{k+1}(\Omega)$  and thus

$$S = \hat{S} + \text{symGrad} v \in \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}(\Omega) + \text{symGrad} \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \subset \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega).$$

For the directness, let  $S = \mathcal{P}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^k T \in \mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega)$  with some  $T \in \mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{Div}, \Omega)$ . Then  $0 = \text{RotRot}^\top S = T$  and thus  $S = 0$ . The assertions about the corresponding Div-spaces follow analogously. Let  $v \in \mathbf{H}_{\Gamma_t}^k(\text{symGrad}, \Omega)$ . Then  $\text{symGrad} v \in \mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega)$  and Theorem 3.1 yields  $\hat{v} := \mathcal{P}_{\text{symGrad},\Gamma_t}^k \text{symGrad} v \in \mathbf{H}_{\Gamma_t}^{k+1}(\Omega)$ . As  $\text{symGrad}(v - \hat{v}) = 0$ , we have  $v - \hat{v} =: r \in \mathbb{R}\mathbb{M}$ , which even vanishes if  $\Gamma_t \neq \emptyset$ . Hence,  $v = \hat{v} + r \in \mathbf{H}_{\Gamma_t}^{k+1}(\Omega)$ .  $\square$

By similar arguments, we also obtain the following (non-standard) versions of Corollaries 3.4 and 3.6.

**Corollary 3.7** (Corollaries 3.4 and 3.6 for nonstandard Sobolev spaces). *Let  $(\Omega, \Gamma_t)$  be an extendable bounded strong Lipschitz pair and let  $k \geq 1$ . Then the bounded regular decompositions*

$$\begin{aligned} \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k, k-1}(\text{RotRot}^\top, \Omega) &= \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega) + \text{symGrad } \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) = R(\mathcal{P}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k-1}) \dot{+} \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega) \\ &= R(\mathcal{P}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k-1}) \dot{+} \text{symGrad } \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \\ &= R(\mathcal{P}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k-1}) \dot{+} \text{symGrad } R(\mathcal{P}_{\text{symGrad}, \Gamma_t}^k) = \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k, k-1}(\text{RotRot}^\top, \Omega) \end{aligned}$$

hold with bounded linear regular decomposition operators

$$\begin{aligned} \mathcal{Q}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k, k-1, 1} &:= \mathcal{P}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k-1} \text{RotRot}^\top : \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k, k-1}(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega), \\ \mathcal{Q}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k, k-1, 0} &:= \mathcal{P}_{\text{symGrad}, \Gamma_t}^k (1 - \mathcal{Q}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k, k-1, 1}) : \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k, k-1}(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \end{aligned}$$

satisfying  $\mathcal{Q}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k, k-1, 1} + \text{symGrad } \mathcal{Q}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k, k-1, 0} = \text{id}_{\mathbf{H}_{\mathbb{S}, \Gamma_t}^{k, k-1}(\text{RotRot}^\top, \Omega)}$ . In particular, weak and strong boundary conditions coincide also for the non-standard Sobolev spaces.

Recall the Hilbert complexes and cohomology groups from Sections 2.6 and 2.7.

**Theorem 3.8** (closed and exact Hilbert complexes for extendable domains). *Let  $(\Omega, \Gamma_t)$  be an extendable bounded strong Lipschitz pair and let  $k \geq 0$ . The domain complexes of linear elasticity*

$$\begin{array}{ccccccccccc} \mathbb{R}M_{\Gamma_t} & \xrightarrow{\mathcal{I}M_{\Gamma_t}} & \mathbf{H}_{\Gamma_t}^{k+1} & \xrightarrow{\text{symGrad}_{\Gamma_t}^k} & \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top) & \xrightarrow{\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}} & \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}) & \xrightarrow{\text{Div}_{\mathbb{S}, \Gamma_t}^k} & \mathbf{H}_{\Gamma_t}^k & \xrightarrow{\mathcal{N}M_{\Gamma_n}} & \mathbb{R}M_{\Gamma_n}, \\ \mathbb{R}M_{\Gamma_t} & \xleftarrow{\mathcal{N}M_{\Gamma_t}} & \mathbf{H}_{\Gamma_t}^k & \xleftarrow{-\text{Div}_{\mathbb{S}, \Gamma_n}^k} & \mathbf{H}_{\mathbb{S}, \Gamma_n}^k(\text{Div}) & \xleftarrow{\text{RotRot}_{\mathbb{S}, \Gamma_n}^{\top, k}} & \mathbf{H}_{\mathbb{S}, \Gamma_n}^k(\text{RotRot}^\top) & \xleftarrow{-\text{symGrad}_{\Gamma_n}^k} & \mathbf{H}_{\Gamma_n}^{k+1} & \xleftarrow{\mathcal{I}M_{\Gamma_n}} & \mathbb{R}M_{\Gamma_n}, \end{array}$$

and, for  $k \geq 1$ ,

$$\begin{array}{ccccccccccc} \mathbb{R}M_{\Gamma_t} & \xrightarrow{\mathcal{I}M_{\Gamma_t}} & \mathbf{H}_{\Gamma_t}^{k+1} & \xrightarrow{\text{symGrad}_{\Gamma_t}^k} & \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k, k-1}(\text{RotRot}^\top) & \xrightarrow{\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k, k-1}} & \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k-1}(\text{Div}) & \xrightarrow{\text{Div}_{\mathbb{S}, \Gamma_t}^{k-1}} & \mathbf{H}_{\Gamma_t}^{k-1} & \xrightarrow{\mathcal{N}M_{\Gamma_n}} & \mathbb{R}M_{\Gamma_n}, \\ \mathbb{R}M_{\Gamma_t} & \xleftarrow{\mathcal{N}M_{\Gamma_t}} & \mathbf{H}_{\Gamma_t}^{k-1} & \xleftarrow{-\text{Div}_{\mathbb{S}, \Gamma_n}^{k-1}} & \mathbf{H}_{\mathbb{S}, \Gamma_n}^{k-1}(\text{Div}) & \xleftarrow{\text{RotRot}_{\mathbb{S}, \Gamma_n}^{\top, k, k-1}} & \mathbf{H}_{\mathbb{S}, \Gamma_n}^{k, k-1}(\text{RotRot}^\top) & \xleftarrow{-\text{symGrad}_{\Gamma_n}^k} & \mathbf{H}_{\Gamma_n}^{k+1} & \xleftarrow{\mathcal{I}M_{\Gamma_n}} & \mathbb{R}M_{\Gamma_n}, \end{array}$$

are exact and closed Hilbert complexes. In particular, all ranges are closed, all cohomology groups (Dirichlet/Neumann fields) are trivial, and the operators from Theorem 3.1 are associated bounded regular potential operators.

### 3.1.2 | General strong Lipschitz domains

Similar to lemma 4.8,<sup>10,11</sup> we get the following.

**Lemma 3.9** (cutting lemma). *Let  $\varphi \in C^\infty(\mathbb{R}^3)$  and let  $k \geq 0$ .*

- (i) *If  $T \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega)$ , then  $\varphi T \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega)$  and  $\text{Div}(\varphi T) = \varphi \text{Div } T + T \text{grad } \varphi$  holds.*
- (ii) *If  $k \geq 1$  and  $S \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k, k-1}(\text{RotRot}^\top, \Omega)$ , then  $\varphi S \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k, k-1}(\text{RotRot}^\top, \Omega)$  and*

$$\text{RotRot}^\top(\varphi S) = \varphi \text{RotRot}^\top S + 2 \text{sym}((\text{spn grad } \varphi) \text{Rot } S) + \Psi(\text{Grad grad } \varphi, S)$$

holds with an algebraic operator  $\Psi$ . In particular, this holds for  $S \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega)$ .

We proceed by showing regular decompositions for the elasticity complexes extending the results of Corollaries 3.4 and 3.7.

**Lemma 3.10** (regular decompositions). *Let  $k \geq 0$ . Then the bounded regular decompositions*

$$\begin{aligned} \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) &= \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega) + \text{RotRot}^\top \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), \\ \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) &= \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}(\Omega) + \text{symGrad} \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega) \end{aligned}$$

and, for  $k \geq 1$ , the nonstandard bounded regular decompositions

$$\mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \subset \mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega) = \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega) + \text{symGrad} \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega)$$

hold with bounded linear regular decomposition operators

$$\begin{aligned} \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,1} &: \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), & \mathcal{Q}_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,1} &: \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) \rightarrow \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega), \\ \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,0} &: \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega), & \mathcal{Q}_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,0} &: \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) \rightarrow \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), \\ \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,k-1,1} &: \mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega), \\ \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,k-1,0} &: \mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega), \end{aligned}$$

satisfying

$$\begin{aligned} \mathcal{Q}_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,1} + \text{RotRot}^\top \mathcal{Q}_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,0} &= \text{id}_{\mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega)}, \\ \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,1} + \text{symGrad} \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,0} &= \text{id}_{\mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega)}, \\ \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,k-1,1} + \text{symGrad} \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,k-1,0} &= \text{id}_{\mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega)}, \quad k \geq 1. \end{aligned}$$

It holds  $\text{Div} \mathcal{Q}_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,1} = \text{Div}_{\mathbb{S},\Gamma_t}^k$  and thus  $\mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{Div}, \Omega)$  is invariant under  $\mathcal{Q}_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,1}$ . Analogously,  $\text{RotRot}^\top \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,1} = \text{RotRot}^\top_{\mathbb{S},\Gamma_t} \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,1}$  and  $\text{RotRot}^\top \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,k-1,1} = \text{RotRot}^\top_{\mathbb{S},\Gamma_t} \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,k-1,1}$  and thus  $\mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega)$  is invariant under  $\mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,1}$  and  $\mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,k-1,1}$ , respectively.

**Corollary 3.11** (weak and strong partial boundary conditions coincide). *Let  $k \geq 0$ . Weak and strong boundary conditions coincide, that is,*

$$\begin{aligned} \mathbf{H}_{\Gamma_t}^k(\text{symGrad}, \Omega) &= \mathbf{H}_{\Gamma_t}^k(\text{symGrad}, \Omega) = \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) = \mathbf{H}_{\Gamma_t}^{k+1}(\Omega), \\ \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) &= \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega), \\ \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) &= \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega), \\ \mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega) &= \mathbf{H}_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega), \quad k \geq 1. \end{aligned}$$

In particular,  $\text{symGrad}_{\Gamma_t}^k = \text{symGrad}_{\Gamma_t}^k$ ,  $\text{RotRot}^\top_{\mathbb{S},\Gamma_t} \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,1} = \text{RotRot}^\top_{\mathbb{S},\Gamma_t} \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,1}$ , and  $\text{Div}_{\mathbb{S},\Gamma_t}^k = \text{Div}_{\mathbb{S},\Gamma_t}^k$ , as well as, for  $k \geq 1$ ,  $\text{RotRot}^\top_{\mathbb{S},\Gamma_t} \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,k-1,1} = \text{RotRot}^\top_{\mathbb{S},\Gamma_t} \mathcal{Q}_{\text{RotRot}^\top_{\mathbb{S},\Gamma_t}}^{k,k-1,1}$ .

*Proof of Lemma 3.10 and Corollary 3.11.* According to references<sup>1,2</sup> and previous works<sup>10,11</sup> (cf. references<sup>12,20,21</sup>), let  $(U_\ell, \varphi_\ell)$  be a partition of unity for  $\Omega$ , that is,

$$\Omega = \bigcup_{\ell=-L}^L \Omega_\ell, \quad \Omega_\ell := \Omega \cap U_\ell, \quad \varphi_\ell \in \mathbf{C}_{\partial U_\ell}^\infty(U_\ell),$$

and  $(\Omega_\ell, \hat{\Gamma}_{t,\ell})$  are extendable bounded strong Lipschitz pairs. Recall  $\Gamma_{t,\ell} := \Gamma_t \cap U_\ell$  and  $\hat{\Gamma}_{t,\ell}$ .

- Let  $k \geq 0$  and let  $T \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega)$ . Then by definition,  $T|_{\Omega_\ell} \in \mathbf{H}_{\mathbb{S}, \Gamma_{t,\ell}}^k(\text{Div}, \Omega_\ell)$ , and we decompose by Corollary 3.4

$$T|_{\Omega_\ell} = T_{\ell,1} + \text{RotRot}^\top S_{\ell,0}$$

with  $T_{\ell,1} := \mathcal{Q}_{\text{Div}\mathbb{S}, \Gamma_{t,\ell}}^{k,1} T|_{\Omega_\ell} \in \mathbf{H}_{\mathbb{S}, \Gamma_{t,\ell}}^{k+1}(\Omega_\ell)$  and  $S_{\ell,0} := \mathcal{Q}_{\text{Div}\mathbb{S}, \Gamma_{t,\ell}}^{k,0} T|_{\Omega_\ell} \in \mathbf{H}_{\mathbb{S}, \Gamma_{t,\ell}}^{k+2}(\Omega_\ell)$ . Lemma 3.9 yields

$$\begin{aligned} \varphi_\ell T|_{\Omega_\ell} &= \varphi_\ell T_{\ell,1} + \varphi_\ell \text{RotRot}^\top S_{\ell,0} \\ &= \overbrace{\varphi_\ell T_{\ell,1} - 2\text{sym}((\text{spn grad } \varphi_\ell)\text{Rot}S_{\ell,0}) - \Psi(\text{Grad grad } \varphi_\ell, S_{\ell,0})} =: T_\ell \\ &\quad + \underbrace{\text{RotRot}^\top(\varphi_\ell S_{\ell,0})}_{=: S_\ell} \end{aligned}$$

with  $T_\ell \in \mathbf{H}_{\mathbb{S}, \hat{\Gamma}_{t,\ell}}^{k+1}(\Omega_\ell)$  and  $S_\ell \in \mathbf{H}_{\mathbb{S}, \hat{\Gamma}_{t,\ell}}^{k+2}(\Omega_\ell)$ . Extending  $T_\ell$  and  $S_\ell$  by zero to  $\Omega$  gives tensor fields  $\tilde{T}_\ell \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega)$  and  $\tilde{S}_\ell \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega)$  as well as

$$\begin{aligned} T &= \sum_{\ell=-L}^L \varphi_\ell T|_{\Omega_\ell} = \sum_{\ell=-L}^L \tilde{T}_\ell + \text{RotRot}^\top \sum_{\ell=-L}^L \tilde{S}_\ell \\ &\in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega) + \text{RotRot}^\top \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) \subset \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega). \end{aligned}$$

As all operations have been linear and continuous, we set

$$\mathcal{Q}_{\text{Div}\mathbb{S}, \Gamma_t}^{k,1} T := \sum_{\ell=-L}^L \tilde{T}_\ell \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega), \quad \mathcal{Q}_{\text{Div}\mathbb{S}, \Gamma_t}^{k,0} T := \sum_{\ell=-L}^L \tilde{S}_\ell \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega).$$

- Let  $k \geq 1$  and let  $S \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega)$ . Then by definition,  $S|_{\Omega_\ell} \in \mathbf{H}_{\mathbb{S}, \Gamma_{t,\ell}}^{k,k-1}(\text{RotRot}^\top, \Omega_\ell)$  and we decompose by Corollary 3.7

$$S|_{\Omega_\ell} = S_{\ell,1} + \text{symGrad } v_{\ell,0}$$

with  $S_{\ell,1} := \mathcal{Q}_{\text{RotRot}^\top \mathbb{S}, \Gamma_{t,\ell}}^{k,k-1,1} S|_{\Omega_\ell} \in \mathbf{H}_{\mathbb{S}, \Gamma_{t,\ell}}^{k+1}(\Omega_\ell)$  and  $v_{\ell,0} := \mathcal{Q}_{\text{RotRot}^\top \mathbb{S}, \Gamma_{t,\ell}}^{k,k-1,0} S|_{\Omega_\ell} \in \mathbf{H}_{\Gamma_{t,\ell}}^{k+1}(\Omega_\ell)$ . Thus,

$$\begin{aligned} \varphi_\ell S|_{\Omega_\ell} &= \varphi_\ell S_{\ell,1} + \varphi_\ell \text{symGrad } v_{\ell,0} \\ &= \underbrace{\varphi_\ell S_{\ell,1} - \text{sym}(v_{\ell,0}(\text{grad } \varphi_\ell)^\top)}_{=: S_\ell} + \underbrace{\text{symGrad}(\varphi_\ell v_{\ell,0})}_{=: v_\ell} \end{aligned} \tag{6}$$

with  $S_\ell \in \mathbf{H}_{\mathbb{S}, \hat{\Gamma}_{t,\ell}}^{k+1}(\Omega_\ell)$  and  $v_\ell \in \mathbf{H}_{\hat{\Gamma}_{t,\ell}}^{k+1}(\Omega_\ell)$ . Extending  $S_\ell$  and  $v_\ell$  by zero to  $\Omega$  gives fields  $\tilde{S}_\ell \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega)$  and  $\tilde{v}_\ell \in \mathbf{H}_{\Gamma_t}^{k+1}(\Omega)$  as well as

$$\begin{aligned} S &= \sum_{\ell=-L}^L \varphi_\ell S|_{\Omega_\ell} = \sum_{\ell=-L}^L \tilde{S}_\ell + \text{symGrad} \sum_{\ell=-L}^L \tilde{v}_\ell \\ &\in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega) + \text{symGrad } \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \subset \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega). \end{aligned}$$

As all operations have been linear and continuous, we set

$$\mathcal{Q}_{\text{RotRot}^\top \mathbb{S}, \Gamma_t}^{k,k-1,1} S := \sum_{\ell=-L}^L \tilde{S}_\ell \in \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega), \quad \mathcal{Q}_{\text{RotRot}^\top \mathbb{S}, \Gamma_t}^{k,k-1,0} S := \sum_{\ell=-L}^L \tilde{v}_\ell \in \mathbf{H}_{\Gamma_t}^{k+1}(\Omega).$$

- Let  $k \geq 0$  and let  $S \in \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega)$ . Then by definition,  $S|_{\Omega_\ell} \in \mathbf{H}_{\mathbb{S},\Gamma_{t,\ell}}^k(\text{RotRot}^\top, \Omega_\ell)$  and we decompose by Corollary 3.4

$$S|_{\Omega_\ell} = S_{\ell,1} + \text{symGrad} v_{\ell,0}$$

with  $S_{\ell,1} := \mathcal{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_{t,\ell}}^\top}^{k,1} S|_{\Omega_\ell} \in \mathbf{H}_{\mathbb{S},\Gamma_{t,\ell}}^{k+2}(\Omega_\ell)$  and  $v_{\ell,0} := \mathcal{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_{t,\ell}}^\top}^{k,0} S|_{\Omega_\ell} \in \mathbf{H}_{\Gamma_{t,\ell}}^{k+1}(\Omega_\ell)$ . Now, we follow the arguments from (6). Note that still only  $S_\ell \in \mathbf{H}_{\mathbb{S},\Gamma_{t,\ell}}^{k+1}(\Omega_\ell)$  holds; that is, we have lost one order of regularity for  $S_\ell$ . Nevertheless, we get

$$S \in \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega) + \text{symGrad} \mathbf{H}_{\Gamma_t}^{k+1}(\Omega),$$

and all operations have been linear and continuous. But this implies by the previous step

$$S \in \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1,k}(\text{RotRot}^\top, \Omega) + \text{symGrad} \mathbf{H}_{\Gamma_t}^{k+1}(\Omega).$$

Again by the previous step, we obtain

$$\begin{aligned} S &\in \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}(\Omega) + \text{symGrad} \mathbf{H}_{\Gamma_t}^{k+2}(\Omega) + \text{symGrad} \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \\ &= \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}(\Omega) + \text{symGrad} \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \subset \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega), \end{aligned}$$

and all operations have been linear and continuous.

It remains to prove  $\mathbf{H}_{\Gamma_t}^k(\text{symGrad}, \Omega) \subset \mathbf{H}_{\Gamma_t}^k(\text{symGrad}, \Omega)$ . Let  $v \in \mathbf{H}_{\Gamma_t}^k(\text{symGrad}, \Omega)$ . Then we have  $\varphi_\ell v \in \mathbf{H}_{\Gamma_{t,\ell}}^k(\text{symGrad}, \Omega_\ell) = \mathbf{H}_{\hat{\Gamma}_{t,\ell}}^k(\text{symGrad}, \Omega_\ell) = \mathbf{H}_{\hat{\Gamma}_{t,\ell}}^{k+1}(\Omega_\ell)$  by Corollary 3.6. Extending  $\varphi_\ell v$  by zero to  $\Omega$  yields vector fields  $v_\ell \in \mathbf{H}_{\Gamma_t}^{k+1}(\Omega)$  and  $v = \sum_\ell \varphi_\ell v = \sum_\ell v_\ell \in \mathbf{H}_{\Gamma_t}^{k+1}(\Omega)$ .  $\square$

### 3.2 | Mini FA-ToolBox

#### 3.2.1 | Zero-order mini FA-ToolBox

Recall Section 2.7 and let  $\varepsilon, \mu$  be admissible. In Section 2.2 (for  $\varepsilon = \mu = \text{id}$ ), we have seen that the densely defined and closed linear operators

$$\begin{aligned} A_0 &= \text{symGrad}_{\Gamma_t} : \mathbf{H}_{\Gamma_t}^1(\Omega) \subset \mathbf{L}^2(\Omega) \rightarrow \mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega), \\ A_1 &= \mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top : \mathbf{H}_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) \subset \mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega) \rightarrow \mathbf{L}_{\mathbb{S},\mu}^2(\Omega), \\ A_2 &= \text{Div}_{\mathbb{S},\Gamma_t} \mu : \mu^{-1} \mathbf{H}_{\mathbb{S},\Gamma_t}(\text{Div}, \Omega) \subset \mathbf{L}_{\mathbb{S},\mu}^2(\Omega) \rightarrow \mathbf{L}^2(\Omega), \\ A_0^* &= -\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon : \varepsilon^{-1} \mathbf{H}_{\mathbb{S},\Gamma_n}(\text{Div}, \Omega) \subset \mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega) \rightarrow \mathbf{L}^2(\Omega), \\ A_1^* &= \varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top : \mathbf{H}_{\mathbb{S},\Gamma_n}(\text{RotRot}^\top, \Omega) \subset \mathbf{L}_{\mathbb{S},\mu}^2(\Omega) \rightarrow \mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega), \\ A_2^* &= -\text{symGrad}_{\Gamma_n} : \mathbf{H}_{\Gamma_n}^1(\Omega) \subset \mathbf{L}^2(\Omega) \rightarrow \mathbf{L}_{\mathbb{S},\mu}^2(\Omega), \end{aligned}$$

where we have used Corollary 3.11, build the long primal and dual elasticity Hilbert complex

$$\begin{array}{ccccccccc} \mathbb{R}\mathbb{M}_{\Gamma_t} & \xrightleftharpoons[A_{-1}^* = \mathbb{R}\mathbb{M}_{\Gamma_t}]{A_{-1} = \mathbb{R}\mathbb{M}_{\Gamma_t}} & \mathbf{L}^2(\Omega) & \xrightleftharpoons[A_0^* = -\text{Div}_{\mathbb{S},\Gamma_n}]{A_0 = \text{symGrad}_{\Gamma_t}} & \mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega) & \xrightleftharpoons[A_1^* = \varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top]{A_1 = \mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top} & \mathbf{L}_{\mathbb{S},\mu}^2(\Omega) & \xrightleftharpoons[A_2^* = -\text{symGrad}_{\Gamma_n}]{A_2 = \text{Div}_{\mathbb{S},\Gamma_t} \mu} & \mathbf{L}^2(\Omega) & \xrightleftharpoons[A_3^* = \mathbb{R}\mathbb{M}_{\Gamma_n}]{A_3 = \mathbb{R}\mathbb{M}_{\Gamma_n}} & \mathbb{R}\mathbb{M}_{\Gamma_n}. \end{array} \quad (7)$$

Compare (5).

**Theorem 3.12** (compact embedding). *The embedding*

$$\mathbf{H}_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) \cap \varepsilon^{-1} \mathbf{H}_{\mathbb{S},\Gamma_n}(\text{Div}, \Omega) \hookrightarrow \mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega)$$

is compact. Moreover, the compactness does not depend on  $\varepsilon$ .



*Proof.* Note that this type of compact embedding is independent of  $\varepsilon$  and  $\mu$  (cf. lemma 5.1<sup>22,23</sup>). So, let  $\varepsilon = \mu = id$ . Lemma 3.10 (for  $k = 0$ ) yields the bounded regular decomposition

$$D(A_0^*) = H_{\mathbb{S},\Gamma_n}(\text{Div}, \Omega) = H_{\mathbb{S},\Gamma_n}^1(\Omega) + \text{RotRot}^\top H_{\mathbb{S},\Gamma_n}^2(\Omega) = H_1^+ + A_1^* H_2^+$$

with  $H_1^+ = H_{\mathbb{S},\Gamma_n}^1(\Omega)$  and  $H_2^+ = H_{\mathbb{S},\Gamma_n}^2(\Omega)$  and  $H_1 = H_2 = L_{\mathbb{S}}^2(\Omega)$ . Rellich's selection theorem and corollary 2.12<sup>10,11</sup> (cf. lemma 2.22<sup>1,2</sup>) yield that  $D(A_1) \cap D(A_0^*) \hookrightarrow H_1$  is compact.  $\square$

*Remark 3.13.* (compact embedding) The embeddings

$$\begin{aligned} D(A_0) \cap D(A_{-1}^*) &= H_{\Gamma_t}^1(\Omega) \hookrightarrow L^2(\Omega) = H_0, \\ D(A_1) \cap D(A_0^*) &= H_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) \cap \varepsilon^{-1} H_{\mathbb{S},\Gamma_n}(\text{Div}, \Omega) \hookrightarrow L_{\mathbb{S},\varepsilon}^2(\Omega) = H_1, \\ D(A_2) \cap D(A_1^*) &= \mu^{-1} H_{\mathbb{S},\Gamma_t}(\text{Div}, \Omega) \cap H_{\mathbb{S},\Gamma_n}(\text{RotRot}^\top, \Omega) \hookrightarrow L_{\mathbb{S},\mu}^2(\Omega) = H_2, \\ D(A_3) \cap D(A_2^*) &= H_{\Gamma_n}^1(\Omega) \hookrightarrow L^2(\Omega) = H_3 \end{aligned}$$

are compact, and the compactness does not depend on  $\varepsilon$  or  $\mu$ .

**Theorem 3.14** (compact elasticity complex). *The long primal and dual elasticity Hilbert complex (7) is compact. In particular, the complex is closed.*

Let us recall the reduced operators

$$\begin{aligned} (A_0)_\perp &= (\text{symGrad}_{\Gamma_t})_\perp : D((\text{symGrad}_{\Gamma_t})_\perp) \subset (\mathbb{R}\mathbb{M}_{\Gamma_t})^{\perp L^2(\Omega)} \rightarrow R(\text{symGrad}_{\Gamma_t}), \\ (A_1)_\perp &= (\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top)_\perp : D((\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top)_\perp) \subset N(\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top)^{\perp L_{\mathbb{S},\varepsilon}^2(\Omega)} \rightarrow R(\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top), \\ (A_2)_\perp &= (\text{Div}_{\mathbb{S},\Gamma_t} \mu)_\perp : D((\text{Div}_{\mathbb{S},\Gamma_t} \mu)_\perp) \subset N(\text{Div}_{\mathbb{S},\Gamma_t} \mu)^{\perp L_{\mathbb{S},\mu}^2(\Omega)} \rightarrow R(\text{Div}_{\mathbb{S},\Gamma_t} \mu), \\ (A_0^*)_\perp &= -(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon)_\perp : D((\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon)_\perp) \subset N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon)^{\perp L_{\mathbb{S},\varepsilon}^2(\Omega)} \rightarrow R(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon), \\ (A_1^*)_\perp &= (\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top)_\perp : D((\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top)_\perp) \subset N(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top)^{\perp L_{\mathbb{S},\mu}^2(\Omega)} \rightarrow R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top), \\ (A_2^*)_\perp &= (\text{symGrad}_{\Gamma_n})_\perp : D((\text{symGrad}_{\Gamma_n})_\perp) \subset (\mathbb{R}\mathbb{M}_{\Gamma_n})^{\perp L^2(\Omega)} \rightarrow R(\text{symGrad}_{\Gamma_n}), \end{aligned}$$

with domains of definition

$$\begin{aligned} D((A_0)_\perp) &= D(\text{symGrad}_{\Gamma_t}) \cap (\mathbb{R}\mathbb{M}_{\Gamma_t})^{\perp L^2(\Omega)}, \\ D((A_1)_\perp) &= D(\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top) \cap N(\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top)^{\perp L_{\mathbb{S},\varepsilon}^2(\Omega)} = D(\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top) \cap R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top), \\ D((A_2)_\perp) &= D(\text{Div}_{\mathbb{S},\Gamma_t} \mu) \cap N(\text{Div}_{\mathbb{S},\Gamma_t} \mu)^{\perp L_{\mathbb{S},\mu}^2(\Omega)} = D(\text{Div}_{\mathbb{S},\Gamma_t} \mu) \cap R(\text{symGrad}_{\Gamma_n}), \\ D((A_0^*)_\perp) &= D(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon) \cap N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon)^{\perp L_{\mathbb{S},\varepsilon}^2(\Omega)} = D(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon) \cap R(\text{symGrad}_{\Gamma_t}), \\ D((A_1^*)_\perp) &= D(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top) \cap N(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top)^{\perp L_{\mathbb{S},\mu}^2(\Omega)} = D(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top) \cap R(\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top), \\ D((A_2^*)_\perp) &= D(\text{symGrad}_{\Gamma_n}) \cap (\mathbb{R}\mathbb{M}_{\Gamma_n})^{\perp L^2(\Omega)}. \end{aligned}$$

Note that  $R(A_n) = R((A_n)_\perp)$  and  $R(A_n^*) = R((A_n^*)_\perp)$  hold. Lemma 2.9<sup>1,2</sup> shows the following:

**Theorem 3.15** (mini FA-ToolBox). *For the zero-order elasticity complex, the following holds:*

- (i) *The ranges  $R(\text{symGrad}_{\Gamma_t})$ ,  $R(\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top)$ , and  $R(\text{Div}_{\mathbb{S},\Gamma_t} \mu)$  are closed.*
- (ii) *The inverse operators  $(\text{symGrad}_{\Gamma_t})_\perp^{-1}$ ,  $(\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top)_\perp^{-1}$ , and  $(\text{Div}_{\mathbb{S},\Gamma_t} \mu)_\perp^{-1}$  are compact.*
- (iii) *The cohomology group of generalized Dirichlet/Neumann tensor fields  $\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega)$  is finite-dimensional. Moreover, the dimension does not depend on  $\varepsilon$ .*

(iv) *The orthonormal Helmholtz-type decompositions*

$$\begin{aligned} L^2_{\mathbb{S},\varepsilon}(\Omega) &= R(\text{symGrad}_{\Gamma_t}) \oplus_{L^2_{\mathbb{S},\varepsilon}(\Omega)} N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon) \\ &= N(\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top) \oplus_{L^2_{\mathbb{S},\varepsilon}(\Omega)} R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top) \\ &= R(\text{symGrad}_{\Gamma_t}) \oplus_{L^2_{\mathbb{S},\varepsilon}(\Omega)} \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) \oplus_{L^2_{\mathbb{S},\varepsilon}(\Omega)} R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top) \end{aligned}$$

hold.

(v) *There exist (optimal)  $c_0, c_1, c_2 > 0$  such that the Friedrichs/Poincaré-type estimates*

$$\begin{aligned} \forall v \in H^1_{\Gamma_t}(\Omega) \cap (\mathbb{R}M_{\Gamma_t})^{\perp L^2(\Omega)} & \quad |v|_{L^2(\Omega)} \leq c_0 |\text{symGrad } v|_{L^2_{\mathbb{S},\varepsilon}(\Omega)}, \\ \forall T \in \varepsilon^{-1} H_{\mathbb{S},\Gamma_n}(\text{Div}, \Omega) \cap R(\text{symGrad}_{\Gamma_t}) & \quad |T|_{L^2_{\mathbb{S},\varepsilon}(\Omega)} \leq c_0 |\text{Div } \varepsilon T|_{L^2(\Omega)}, \\ \forall S \in H_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) \cap R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top) & \quad |S|_{L^2_{\mathbb{S},\varepsilon}(\Omega)} \leq c_1 |\mu^{-1} \text{RotRot}^\top S|_{L^2_{\mathbb{S},\mu}(\Omega)}, \\ \forall S \in H_{\mathbb{S},\Gamma_n}(\text{RotRot}^\top, \Omega) \cap R(\mu^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^\top) & \quad |S|_{L^2_{\mathbb{S},\mu}(\Omega)} \leq c_1 |\varepsilon^{-1} \text{RotRot}^\top S|_{L^2_{\mathbb{S},\varepsilon}(\Omega)}, \\ \forall T \in \mu^{-1} H_{\mathbb{S},\Gamma_t}(\text{Div}, \Omega) \cap R(\text{symGrad}_{\Gamma_n}) & \quad |T|_{L^2_{\mathbb{S},\mu}(\Omega)} \leq c_2 |\text{Div } \mu T|_{L^2(\Omega)}, \\ \forall v \in H^1_{\Gamma_n}(\Omega) \cap (\mathbb{R}M_{\Gamma_n})^{\perp L^2(\Omega)} & \quad |v|_{L^2(\Omega)} \leq c_2 |\text{symGrad } v|_{L^2_{\mathbb{S},\mu}(\Omega)} \end{aligned}$$

hold.

(vi) *For all  $S \in H_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) \cap \varepsilon^{-1} H_{\mathbb{S},\Gamma_n}(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega)^{\perp L^2_{\mathbb{S},\varepsilon}(\Omega)}$ , it holds*

$$|S|_{L^2_{\mathbb{S},\varepsilon}(\Omega)}^2 \leq c_1^2 |\mu^{-1} \text{RotRot}^\top S|_{L^2_{\mathbb{S},\mu}(\Omega)}^2 + c_0^2 |\text{Div } \varepsilon S|_{L^2(\Omega)}^2.$$

(vii)  $\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) = \{0\}$ , if  $(\Omega, \Gamma_t)$  is extendable.

### 3.2.2 | Higher order mini FA-ToolBox

For simplicity, let  $\varepsilon = \mu = \text{id}$ . From Section 2.6, we recall the densely defined and closed higher Sobolev order operators

$$\begin{aligned} \text{symGrad}_{\Gamma_t}^k &: H_{\Gamma_t}^{k+1}(\Omega) \subset H_{\Gamma_t}^k(\Omega) \rightarrow H_{\mathbb{S},\Gamma_t}^k(\Omega), \\ \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k} &: H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \subset H_{\mathbb{S},\Gamma_t}^k(\Omega) \rightarrow H_{\mathbb{S},\Gamma_t}^k(\Omega), \\ \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k,k-1} &: H_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega) \subset H_{\mathbb{S},\Gamma_t}^k(\Omega) \rightarrow H_{\mathbb{S},\Gamma_t}^{k-1}(\Omega), \quad k \geq 1, \\ \text{Div}_{\mathbb{S},\Gamma_t}^k &: H_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) \subset H_{\mathbb{S},\Gamma_t}^k(\Omega) \rightarrow H_{\Gamma_t}^k(\Omega), \end{aligned} \tag{8}$$

building the long elasticity Hilbert complexes

$$\mathbb{R}M_{\Gamma_t} \xrightarrow{\mathbb{R}M_{\Gamma_t}} H_{\Gamma_t}^k(\Omega) \xrightarrow{\text{symGrad}_{\Gamma_t}^k} H_{\mathbb{S},\Gamma_t}^k(\Omega) \xrightarrow{\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}} H_{\mathbb{S},\Gamma_t}^k(\Omega) \xrightarrow{\text{Div}_{\mathbb{S},\Gamma_t}^k} H_{\Gamma_t}^k(\Omega) \xrightarrow{\mathbb{R}M_{\Gamma_n}} \mathbb{R}M_{\Gamma_n}, \quad k \geq 0, \tag{9}$$

$$\mathbb{R}M_{\Gamma_t} \xrightarrow{\mathbb{R}M_{\Gamma_t}} H_{\Gamma_t}^k(\Omega) \xrightarrow{\text{symGrad}_{\Gamma_t}^k} H_{\mathbb{S},\Gamma_t}^k(\Omega) \xrightarrow{\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k,k-1}} H_{\mathbb{S},\Gamma_t}^{k-1}(\Omega) \xrightarrow{\text{Div}_{\mathbb{S},\Gamma_t}^{k-1}} H_{\Gamma_t}^{k-1}(\Omega) \xrightarrow{\mathbb{R}M_{\Gamma_n}} \mathbb{R}M_{\Gamma_n}, \quad k \geq 1. \tag{10}$$

We start with regular representations implied by Lemma 3.10 and Corollary 3.11.

**Theorem 3.16** (regular representations and closed ranges). *Let  $k \geq 0$ . Then the regular potential representations*

$$\begin{aligned}
 R(\text{symGrad}_{\Gamma_t}^k) &= \text{symGrad } H_{\Gamma_t}^k(\text{symGrad}, \Omega) = \text{symGrad } H_{\Gamma_t}^{k+1}(\Omega) \\
 &= H_{\mathbb{S}, \Gamma_t}^k(\Omega) \cap R(\text{symGrad}_{\Gamma_t}) \\
 &= H_{\mathbb{S}, \Gamma_t}^k(\Omega) \cap H_{\mathbb{S}, \Gamma_t, 0}(\text{RotRot}^\top, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_t, \Gamma_n, \epsilon}(\Omega)^{\perp_{L^2_{\mathbb{S}}(\Omega)}} \\
 &= H_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_t, \Gamma_n, \epsilon}(\Omega)^{\perp_{L^2_{\mathbb{S}}(\Omega)}}, \\
 R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k}) &= R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}) = \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega) = \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) \\
 &= \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega) \\
 &= H_{\mathbb{S}, \Gamma_t}^k(\Omega) \cap R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top) \\
 &= H_{\mathbb{S}, \Gamma_t}^k(\Omega) \cap H_{\mathbb{S}, \Gamma_t, 0}(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_n, \Gamma_t, \epsilon}(\Omega)^{\perp_{L^2_{\mathbb{S}}(\Omega)}}, \\
 &= H_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_n, \Gamma_t, \epsilon}(\Omega)^{\perp_{L^2_{\mathbb{S}}(\Omega)}}, \\
 R(\text{Div}_{\mathbb{S}, \Gamma_t}^k) &= \text{Div} H_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega) = \text{Div} H_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega) \\
 &= H_{\Gamma_t}^k(\Omega) \cap R(\text{Div}_{\mathbb{S}, \Gamma_t}) = H_{\Gamma_t}^k(\Omega) \cap (\mathbb{RM}_{\Gamma_n})^{\perp_{L^2(\Omega)}}
 \end{aligned}$$

hold. In particular, the latter spaces are closed subspaces of  $H_{\mathbb{S}}^k(\Omega)$  and  $H^k(\Omega)$ , respectively, and all ranges of the higher Sobolev order operators in (8) are closed. Moreover, the long elasticity Hilbert complexes (9) and (10) are closed.

Note that in Theorem 3.16 we claim nothing about bounded regular potential operators, leaving the question of bounded potentials to the next sections (cf. Theorem 3.24).

*Proof of Theorem 3.16.* We only show the representations for  $R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})$ . The others follow analogously, but simpler. By Lemma 3.10 and Corollary 3.11, we have

$$\begin{aligned}
 \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) &\subset \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega) = R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k}) \\
 &\subset \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega) = R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}) = \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega).
 \end{aligned}$$

In particular,

$$R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}) = \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega) = \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega). \tag{11}$$

Moreover,

$$\begin{aligned}
 R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}) &\subset H_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_n, \Gamma_t, \epsilon}(\Omega)^{\perp_{L^2_{\mathbb{S}}(\Omega)}} \\
 &= H_{\mathbb{S}, \Gamma_t}^k(\Omega) \cap H_{\mathbb{S}, \Gamma_t, 0}(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_n, \Gamma_t, \epsilon}(\Omega)^{\perp_{L^2_{\mathbb{S}}(\Omega)}} = H_{\mathbb{S}, \Gamma_t}^k(\Omega) \cap R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top),
 \end{aligned}$$

since by Theorem 3.15 (iv)

$$R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top) = H_{\mathbb{S}, \Gamma_t, 0}(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_n, \Gamma_t, \epsilon}(\Omega)^{\perp_{L^2_{\mathbb{S}}(\Omega)}}. \tag{12}$$

Thus, it remains to show

$$H_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_n, \Gamma_t, \epsilon}(\Omega)^{\perp_{L^2_{\mathbb{S}}(\Omega)}} \subset \text{RotRot}^\top H_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega), \quad k \geq 1.$$

For this, let  $k \geq 1$  and  $T \in H_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_n, \Gamma_t, \epsilon}(\Omega)^{\perp_{H^2_{\mathbb{S}}(\Omega)}}$ . By (12) and (11), we have

$$T \in R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top}) = \text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top} H_{\mathbb{S}, \Gamma_t}^2(\Omega),$$

and hence, there is  $S_1 \in H_{\mathbb{S}, \Gamma_t}^2(\Omega)$  such that  $\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top} S_1 = T$ . We see  $S_1 \in H_{\mathbb{S}, \Gamma_t}^2(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top}, \Omega)$  resp.  $S_1 \in H_{\mathbb{S}, \Gamma_t}^1(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top}, \Omega)$  if  $k = 1$ . Hence, we are done for  $k = 1$  and  $k = 2$ . For  $k \geq 2$ , we have  $T \in \text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top} H_{\mathbb{S}, \Gamma_t}^2(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top}, \Omega) = \text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top} H_{\mathbb{S}, \Gamma_t}^4(\Omega)$  by (11). Thus, there is  $S_2 \in H_{\mathbb{S}, \Gamma_t}^4(\Omega)$  such that  $\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top} S_2 = T$ . Then  $S_2 \in H_{\mathbb{S}, \Gamma_t}^4(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top}, \Omega)$  resp.  $S_2 \in H_{\mathbb{S}, \Gamma_t}^3(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top}, \Omega)$  if  $k = 3$ , and we are done for  $k = 3$  and  $k = 4$ . After finitely many steps, we observe that  $T$  belongs to  $\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top} H_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top}, \Omega)$ , finishing the proof.  $\square$

The reduced operators corresponding to (8) are

$$\begin{aligned} (\text{symGrad}_{\Gamma_t}^k)_{\perp} : D((\text{symGrad}_{\Gamma_t}^k)_{\perp}) \subset (\mathbb{R}\mathbb{M}_{\Gamma_t})^{\perp_{H^k(\Omega)}} &\rightarrow R(\text{symGrad}_{\Gamma_t}^k), \\ (\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})_{\perp} : D((\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})_{\perp}) \subset N(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})^{\perp_{H^k(\Omega)}} &\rightarrow R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}), \\ (\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k, k-1})_{\perp} : D((\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k, k-1})_{\perp}) \subset N(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})^{\perp_{H^k(\Omega)}} &\rightarrow R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k-1}), \quad k \geq 1, \\ (\text{Div}_{\mathbb{S}, \Gamma_t}^k)_{\perp} : D((\text{Div}_{\mathbb{S}, \Gamma_t}^k)_{\perp}) \subset N(\text{Div}_{\mathbb{S}, \Gamma_t}^k)^{\perp_{H^k(\Omega)}} &\rightarrow R(\text{Div}_{\mathbb{S}, \Gamma_t}^k) \end{aligned}$$

with domains of definition

$$\begin{aligned} D((\text{symGrad}_{\Gamma_t}^k)_{\perp}) &= D(\text{symGrad}_{\Gamma_t}^k) \cap (\mathbb{R}\mathbb{M}_{\Gamma_t})^{\perp_{H^k(\Omega)}}, \\ D((\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})_{\perp}) &= D(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}) \cap N(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})^{\perp_{H^k(\Omega)}}, \\ D((\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k, k-1})_{\perp}) &= D(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k, k-1}) \cap N(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})^{\perp_{H^k(\Omega)}}, \quad k \geq 1, \\ D((\text{Div}_{\mathbb{S}, \Gamma_t}^k)_{\perp}) &= D(\text{Div}_{\mathbb{S}, \Gamma_t}^k) \cap N(\text{Div}_{\mathbb{S}, \Gamma_t}^k)^{\perp_{H^k(\Omega)}}. \end{aligned}$$

Lemma 2.1<sup>1,2</sup> and Theorem 3.16 yield:

**Theorem 3.17** (closed ranges and bounded inverse operators). *Let  $k \geq 0$ . Then:*

(i)  $R(\text{symGrad}_{\Gamma_t}^k) = R((\text{symGrad}_{\Gamma_t}^k)_{\perp})$  are closed and, equivalently, the inverse operator

$$\begin{aligned} (\text{symGrad}_{\Gamma_t}^k)_{\perp}^{-1} : R(\text{symGrad}_{\Gamma_t}^k) &\rightarrow D((\text{symGrad}_{\Gamma_t}^k)_{\perp}) \\ \text{resp. } (\text{symGrad}_{\Gamma_t}^k)_{\perp}^{-1} : R(\text{symGrad}_{\Gamma_t}^k) &\rightarrow D(\text{symGrad}_{\Gamma_t}^k) \end{aligned}$$

is bounded. Equivalently, there is  $c > 0$  such that for all  $v \in D((\text{symGrad}_{\Gamma_t}^k)_{\perp})$

$$|v|_{H^k(\Omega)} \leq c |\text{symGrad } v|_{H_{\mathbb{S}}^k(\Omega)}.$$

(ii)  $R((\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})_{\perp}) = R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}) = R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k}) = R((\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k})_{\perp})$  are closed and, equivalently, the inverse operators

$$\begin{aligned}
 & (\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})_{\perp}^{-1} : R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}) \rightarrow D((\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})_{\perp}) \\
 \text{resp.} \quad & (\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})_{\perp}^{-1} : R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}) \rightarrow D(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}), \\
 & (\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k})_{\perp}^{-1} : R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}) \rightarrow D((\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k})_{\perp}) \\
 \text{resp.} \quad & (\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k})_{\perp}^{-1} : R(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}) \rightarrow D(\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k})
 \end{aligned}$$

are bounded. Equivalently, there is  $c > 0$  such that for all  $S \in D((\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})_{\perp})$  resp.  $S \in D((\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k})_{\perp})$

$$|S|_{\mathbb{H}_{\mathbb{S}}^k(\Omega)} \leq c|\text{RotRot}^{\top} S|_{\mathbb{H}_{\mathbb{S}}^k(\Omega)} \text{ resp. } |S|_{\mathbb{H}_{\mathbb{S}}^{k+1}(\Omega)} \leq c|\text{RotRot}^{\top} S|_{\mathbb{H}_{\mathbb{S}}^k(\Omega)}.$$

(iii)  $R(\text{Div}_{\mathbb{S}, \Gamma_t}^k) = R((\text{Div}_{\mathbb{S}, \Gamma_t}^k)_{\perp})$  are closed and, equivalently, the inverse operator

$$\begin{aligned}
 & (\text{Div}_{\mathbb{S}, \Gamma_t}^k)_{\perp}^{-1} : R(\text{Div}_{\mathbb{S}, \Gamma_t}^k) \rightarrow D((\text{Div}_{\mathbb{S}, \Gamma_t}^k)_{\perp}) \\
 \text{resp.} \quad & (\text{Div}_{\mathbb{S}, \Gamma_t}^k)_{\perp}^{-1} : R(\text{Div}_{\mathbb{S}, \Gamma_t}^k) \rightarrow D(\text{Div}_{\mathbb{S}, \Gamma_t}^k)
 \end{aligned}$$

is bounded. Equivalently, there is  $c > 0$  such that for all  $T \in D((\text{Div}_{\mathbb{S}, \Gamma_t}^k)_{\perp})$

$$|T|_{\mathbb{H}_{\mathbb{S}}^k(\Omega)} \leq c|\text{Div} T|_{\mathbb{H}^k(\Omega)}.$$

**Lemma 3.18.** (Schwarz' lemma). Let  $0 \leq |\alpha| \leq k$ .

- (i) For  $S \in \mathbb{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^{\top}, \Omega)$  resp.  $S \in \mathbb{H}_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^{\top}, \Omega)$ , it holds  $\partial^{\alpha} S \in \mathbb{H}_{\mathbb{S}, \Gamma_t}(\text{RotRot}^{\top}, \Omega)$  resp.  $\partial^{\alpha} S \in \mathbb{H}_{\mathbb{S}, \Gamma_t}^{1, 0}(\text{RotRot}^{\top}, \Omega)$  and  $\text{RotRot}^{\top} \partial^{\alpha} S = \partial^{\alpha} \text{RotRot}^{\top} S$ .
- (ii) For  $T \in \mathbb{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega)$ , it holds  $\partial^{\alpha} T \in \mathbb{H}_{\mathbb{S}, \Gamma_t}(\text{Div}, \Omega)$  and  $\text{Div} \partial^{\alpha} T = \partial^{\alpha} \text{Div} T$ .

*Proof.* Let  $S \in \mathbb{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^{\top}, \Omega)$ . For  $\Phi \in C_{\Gamma_n}^{\infty}(\Omega)$ , we have

$$\begin{aligned}
 \langle \partial^{\alpha} S, \text{RotRot}^{\top} \Phi \rangle_{L_{\mathbb{S}}^2(\Omega)} &= (-1)^{|\alpha|} \langle S, \text{RotRot}^{\top} \partial^{\alpha} \Phi \rangle_{L_{\mathbb{S}}^2(\Omega)} \\
 &= (-1)^{|\alpha|} \langle \text{RotRot}^{\top} S, \partial^{\alpha} \Phi \rangle_{L_{\mathbb{S}}^2(\Omega)} = \langle \partial^{\alpha} \text{RotRot}^{\top} S, \Phi \rangle_{L_{\mathbb{S}}^2(\Omega)}
 \end{aligned}$$

as  $S \in \mathbb{H}_{\mathbb{S}, \Gamma_t}^k(\Omega) \cap \mathbb{H}_{\mathbb{S}, \Gamma_t}(\text{RotRot}^{\top}, \Omega)$  and  $\text{RotRot}^{\top} S \in \mathbb{H}_{\mathbb{S}, \Gamma_t}^k(\Omega)$ . Hence,

$$\partial^{\alpha} S \in \mathbf{H}_{\mathbb{S}, \Gamma_t}(\text{RotRot}^{\top}, \Omega) = \mathbb{H}_{\mathbb{S}, \Gamma_t}(\text{RotRot}^{\top}, \Omega)$$

by Corollary 3.11 and  $\text{RotRot}^{\top} \partial^{\alpha} S = \partial^{\alpha} \text{RotRot}^{\top} S$ . The other assertions follow analogously. □

**Theorem 3.19** (compact embedding). Let  $k \geq 0$ . Then the embedding

$$\mathbb{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^{\top}, \Omega) \cap \mathbb{H}_{\mathbb{S}, \Gamma_n}^k(\text{Div}, \Omega) \hookrightarrow \mathbb{H}_{\mathbb{S}, \Gamma}^k(\Omega)$$

is compact.

*Proof.* We follow in close lines the proof of theorem 4.11<sup>10,11</sup> (cf. theorem 4.16<sup>1,2</sup>) using induction. The case  $k = 0$  is given by Theorem 3.12. Let  $k \geq 1$  and let  $(S_{\ell})$  be a bounded sequence in  $\mathbb{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^{\top}, \Omega) \cap \mathbb{H}_{\mathbb{S}, \Gamma_n}^k(\text{Div}, \Omega)$ . Note that

$$\mathbb{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^{\top}, \Omega) \cap \mathbb{H}_{\mathbb{S}, \Gamma_n}^k(\text{Div}, \Omega) \subset \mathbb{H}_{\mathbb{S}, \Gamma_t}^k(\Omega) \cap \mathbb{H}_{\mathbb{S}, \Gamma_n}^k(\Omega) = \mathbb{H}_{\mathbb{S}, \Gamma}^k(\Omega).$$

By assumption and w.l.o.g., we have that  $(S_\ell)$  is a Cauchy sequence in  $H_{\mathbb{S},\Gamma}^{k-1}(\Omega)$ . Moreover, for all  $|\alpha| = k$ , we have  $\partial^\alpha S_\ell \in H_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) \cap H_{\mathbb{S},\Gamma_n}(\text{Div}, \Omega)$  with  $\text{RotRot}^\top \partial^\alpha S_\ell = \partial^\alpha \text{RotRot}^\top S_\ell$  and  $\text{Div} \partial^\alpha S_\ell = \partial^\alpha \text{Div} S_\ell$  by Lemma 3.18. Hence,  $(\partial^\alpha S_\ell)$  is a bounded sequence in the zero-order space  $H_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) \cap H_{\mathbb{S},\Gamma_n}(\text{Div}, \Omega)$ . Thus, w.l.o.g.  $(\partial^\alpha S_\ell)$  is a Cauchy sequence in  $L^2_{\mathbb{S}}(\Omega)$  by Theorem 3.12. Finally,  $(S_\ell)$  is a Cauchy sequence in  $H_{\mathbb{S},\Gamma}^k(\Omega)$ , finishing the proof.  $\square$

**Remark 3.20** (compact embedding) For  $k \geq 1$  (cf. remark 4.12<sup>10,11</sup>), there is another and slightly more general proof using a variant of lemma 2.22.<sup>1,2</sup>

For this, let  $(S_\ell)$  be a bounded sequence in  $H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \cap H_{\mathbb{S},\Gamma_n}^k(\text{Div}, \Omega)$ . In particular,  $(S_\ell)$  is bounded in  $H_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega) \cap H_{\mathbb{S},\Gamma_n}^k(\text{Div}, \Omega)$ . According to Lemma 3.10, we decompose  $S_\ell = T_\ell + \text{symGrad} v_\ell$  with  $T_\ell \in H_{\mathbb{S},\Gamma_t}^{k+1}(\Omega)$  and  $v_\ell \in H_{\Gamma_t}^{k+1}(\Omega)$ . By the boundedness of the regular decomposition operators,  $(T_\ell)$  and  $(v_\ell)$  are bounded in  $H_{\mathbb{S},\Gamma_t}^{k+1}(\Omega)$  and  $H_{\Gamma_t}^{k+1}(\Omega)$ , respectively. W.l.o.g.  $(T_\ell)$  and  $(v_\ell)$  converge in  $H_{\mathbb{S},\Gamma_t}^k(\Omega)$  and  $H_{\Gamma_t}^k(\Omega)$ , respectively. For all  $0 \leq |\alpha| \leq k$ , Lemma 3.18 yields  $(\partial^\alpha S_\ell) \subset H_{\mathbb{S},\Gamma_n}(\text{Div}, \Omega)$  and  $\text{Div} \partial^\alpha T = \partial^\alpha \text{Div} T$ . With  $S_{\ell,l} := S_\ell - S_l$ ,  $T_{\ell,l} := T_\ell - T_l$ , and  $v_{\ell,l} := v_\ell - v_l$ , we get

$$\begin{aligned} |S_{\ell,l}|_{H_{\mathbb{S}}^k(\Omega)}^2 &= \langle S_{\ell,l}, T_{\ell,l} \rangle_{H_{\mathbb{S}}^k(\Omega)} + \langle S_{\ell,l}, \text{symGrad} v_{\ell,l} \rangle_{H_{\mathbb{S}}^k(\Omega)} \\ &= \langle S_{\ell,l}, T_{\ell,l} \rangle_{H_{\mathbb{S}}^k(\Omega)} - \langle \text{Div} S_{\ell,l}, v_{\ell,l} \rangle_{H^k(\Omega)} \leq c \left( |T_{\ell,l}|_{H_{\mathbb{S}}^k(\Omega)} + |v_{\ell,l}|_{H^k(\Omega)} \right) \rightarrow 0. \end{aligned}$$

The latter remark shows immediately:

**Theorem 3.21** (compact embedding). *Let  $k \geq 1$ . Then the embedding*

$$H_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega) \cap H_{\mathbb{S},\Gamma_n}^k(\text{Div}, \Omega) \hookrightarrow H_{\mathbb{S},\Gamma}^k(\Omega)$$

is compact.

**Theorem 3.22** (Friedrichs/Poincaré-type estimate). *There exists  $\hat{c}_k > 0$  such that for all  $S$  in  $H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \cap H_{\mathbb{S},\Gamma_n}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\text{id}}(\Omega)^{\perp L^2_{\mathbb{S}}(\Omega)}$*

$$|S|_{H_{\mathbb{S}}^k(\Omega)} \leq \hat{c}_k \left( |\text{RotRot}^\top S|_{H_{\mathbb{S}}^k(\Omega)} + |\text{Div} S|_{H^k(\Omega)} \right).$$

The condition  $\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\text{id}}(\Omega)^{\perp L^2_{\mathbb{S}}(\Omega)}$  can be replaced by the weaker conditions  $\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\text{id}}^k(\Omega)^{\perp L^2_{\mathbb{S}}(\Omega)}$  or  $\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\text{id}}^k(\Omega)^{\perp H^k(\Omega)}$ . In particular, it holds

$$\begin{aligned} \forall S \in H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \cap R(\text{RotRot}^\top_{\mathbb{S},\Gamma_n}) & \quad |S|_{H_{\mathbb{S}}^k(\Omega)} \leq \hat{c}_k |\text{RotRot}^\top S|_{H_{\mathbb{S}}^k(\Omega)}, \\ \forall S \in H_{\mathbb{S},\Gamma_n}^k(\text{Div}, \Omega) \cap R(\text{symGrad}^k_{\Gamma_t}) & \quad |S|_{H_{\mathbb{S}}^k(\Omega)} \leq \hat{c}_k |\text{Div} S|_{H^k(\Omega)} \end{aligned}$$

with

$$\begin{aligned} R(\text{RotRot}^\top_{\mathbb{S},\Gamma_n})^{k,k+1,k} &= R(\text{RotRot}^\top_{\mathbb{S},\Gamma_n})^{k,k} = H_{\mathbb{S},\Gamma_n,0}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\text{id}}(\Omega)^{\perp L^2_{\mathbb{S}}(\Omega)}, \\ R(\text{symGrad}^k_{\Gamma_t}) &= H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\text{id}}(\Omega)^{\perp L^2_{\mathbb{S}}(\Omega)}. \end{aligned}$$

Analogously, for  $k \geq 1$ , there exists  $\hat{c}_{k,k-1} > 0$  such that

$$|S|_{H_{\mathbb{S}}^k(\Omega)} \leq \hat{c}_{k,k-1} \left( |\text{RotRot}^\top S|_{H_{\mathbb{S}}^{k-1}(\Omega)} + |\text{Div} S|_{H^k(\Omega)} \right)$$

for all  $S$  in  $H_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega) \cap H_{\mathbb{S},\Gamma_n}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\text{id}}(\Omega)^{\perp L^2_{\mathbb{S}}(\Omega)}$ . Moreover,

$$\forall S \in H_{\mathbb{S},\Gamma_t}^{k,k-1}(\text{RotRot}^\top, \Omega) \cap R(\text{RotRot}^\top_{\mathbb{S},\Gamma_n})^{k,k} \quad |S|_{H_{\mathbb{S}}^k(\Omega)} \leq \hat{c}_{k,k-1} |\text{RotRot}^\top S|_{H_{\mathbb{S}}^{k-1}(\Omega)}.$$

*Proof.* We follow the proof of theorem 4.17.<sup>1,2</sup> To show the first estimate, we use a standard strategy and assume the contrary. Then there is a sequence

$$(S_\ell) \subset H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \cap H_{\mathbb{S},\Gamma_n}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\text{id}}(\Omega) \stackrel{\perp}{L^2_{\mathbb{S}}(\Omega)}$$

with  $|S_\ell|_{H_{\mathbb{S}}^k(\Omega)} = 1$  and  $|\text{RotRot}^\top S_\ell|_{H_{\mathbb{S}}^k(\Omega)} + |\text{Div} S_\ell|_{H^k(\Omega)} \rightarrow 0$ . Hence, we may assume that  $S_\ell$  converges weakly to some  $S$  in  $H_{\mathbb{S}}^k(\Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\text{id}}(\Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\text{id}}(\Omega) \stackrel{\perp}{L^2_{\mathbb{S}}(\Omega)}$ . Thus,  $S = 0$ . By Theorem 3.19,  $(S_\ell)$  converges strongly to 0 in  $H_{\mathbb{S}}^k(\Omega)$ , in contradiction to  $|S_\ell|_{H^k(\Omega)} = 1$ . The other estimates follow analogously resp. with Theorem 3.16 by restriction.  $\square$

*Remark 3.23* (Friedrichs/Poincaré/Korn-type estimate). Similar to Theorem 3.22 and by Rellich's selection theorem, there exists  $c > 0$  such that for all  $v \in H_{\Gamma_t}^{k+1}(\Omega) \cap (\mathbb{R}\mathbb{M}_{\Gamma_t})^{\perp L^2(\Omega)}$

$$|v|_{H^k(\Omega)} \leq c |\text{symGrad} v|_{H_{\mathbb{S}}^k(\Omega)}.$$

As in Theorem 3.17,  $(\mathbb{R}\mathbb{M}_{\Gamma_t})^{\perp L^2(\Omega)}$  can be replaced by  $(\mathbb{R}\mathbb{M}_{\Gamma_t})^{\perp H_{\Gamma_t}^k(\Omega)}$ .

### 3.3 | Regular potentials and decompositions II

Let  $k \geq 0$ . According to Theorem 3.17, the inverses of the reduced operators

$$\begin{aligned} (\text{symGrad}_{\Gamma_t}^k)_\perp^{-1} &: R(\text{symGrad}_{\Gamma_t}^k) \rightarrow D(\text{symGrad}_{\Gamma_t}^k) = H_{\Gamma_t}^{k+1}(\Omega), \\ (\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k})_\perp^{-1} &: R(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}) \rightarrow D(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}) = H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega), \\ (\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k})_\perp^{-1} &: R(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k}) \rightarrow D(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k}) = H_{\mathbb{S},\Gamma_t}^{k+1,k}(\text{RotRot}^\top, \Omega), \\ (\text{Div}_{\mathbb{S},\Gamma_t}^k)_\perp^{-1} &: R(\text{Div}_{\mathbb{S},\Gamma_t}^k) \rightarrow D(\text{Div}_{\mathbb{S},\Gamma_t}^k) = H_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) \end{aligned}$$

are bounded and we recall the bounded linear regular decomposition operators

$$\begin{aligned} \mathcal{Q}_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,1} &: H_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) \rightarrow H_{\mathbb{S},\Gamma_t}^{k+1}(\Omega), & \mathcal{Q}_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,0} &: H_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega) \rightarrow H_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), \\ \mathcal{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1} &: H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow H_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), & \mathcal{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,0} &: H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow H_{\Gamma_t}^{k+1}(\Omega), \\ \mathcal{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k,1} &: H_{\mathbb{S},\Gamma_t}^{k+1,k}(\text{RotRot}^\top, \Omega) \rightarrow H_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), & \mathcal{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k,0} &: H_{\mathbb{S},\Gamma_t}^{k+1,k}(\text{RotRot}^\top, \Omega) \rightarrow H_{\Gamma_t}^{k+2}(\Omega) \end{aligned}$$

from Lemma 3.10. Similar to theorems 4.18 and 5.2<sup>1,2</sup> (cf. lemma 2.22 and theorem 2.23<sup>1,2</sup>), we obtain the following sequence of results:

**Theorem 3.24** (bounded regular potentials from bounded regular decompositions). *For  $k \geq 0$ , there exist bounded linear regular potential operators*

$$\begin{aligned} \mathcal{P}_{\text{symGrad}_{\Gamma_t}}^k &:= (\text{symGrad}_{\Gamma_t}^k)_\perp^{-1} : H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\epsilon}(\Omega) \stackrel{\perp}{L^2_{\mathbb{S},\epsilon}(\Omega)} \rightarrow H_{\Gamma_t}^{k+1}(\Omega), \\ \mathcal{P}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^k &:= \mathcal{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1} (\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k})_\perp^{-1} : H_{\mathbb{S},\Gamma_t,0}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_n,\Gamma_t,\epsilon}(\Omega) \stackrel{\perp}{L^2_{\mathbb{S}}(\Omega)} \rightarrow H_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), \\ \mathcal{P}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k} &:= \mathcal{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k,1} (\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k})_\perp^{-1} : H_{\mathbb{S},\Gamma_t,0}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S},\Gamma_n,\Gamma_t,\epsilon}(\Omega) \stackrel{\perp}{L^2_{\mathbb{S}}(\Omega)} \rightarrow H_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), \\ \mathcal{P}_{\text{Div}_{\mathbb{S},\Gamma_t}}^k &:= \mathcal{Q}_{\text{Div}_{\mathbb{S},\Gamma_t}}^{k,1} (\text{Div}_{\mathbb{S},\Gamma_t}^k)_\perp^{-1} : H_{\Gamma_t}^k(\Omega) \cap (\mathbb{R}\mathbb{M}_{\Gamma_n})^{\perp L^2(\Omega)} \rightarrow H_{\mathbb{S},\Gamma_t}^{k+1}(\Omega), \end{aligned}$$

such that

$$\begin{aligned} \text{symGrad} \mathcal{P}_{\text{symGrad}, \Gamma_t}^k &= \text{id} |_{\mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_t, \Gamma_n, \varepsilon}^k(\Omega)} \stackrel{1,2}{\mathbb{S}, \varepsilon}(\Omega)}, \\ \text{RotRot}^\top \mathcal{P}_{\text{RotRot}^\top, \Gamma_t}^{k+1, k} &= \text{RotRot}^\top \mathcal{P}_{\text{RotRot}^\top, \Gamma_t}^k = \text{id} |_{\mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega) \cap \mathcal{H}_{\mathbb{S}, \Gamma_t, \Gamma_t, \varepsilon}^k(\Omega)} \stackrel{1,2}{\mathbb{S}}(\Omega)}, \\ \text{Div} \mathcal{P}_{\text{Div}, \Gamma_t}^k &= \text{id} |_{\mathbf{H}_{\Gamma_t}^k(\Omega) \cap (\mathbb{R}\mathbb{M}_{\Gamma_n})} \stackrel{1,2}{L^2(\Omega)}. \end{aligned}$$

In particular, all potentials in Theorem 3.16 can be chosen such that they depend continuously on the data.  $\mathcal{P}_{\text{symGrad}, \Gamma_t}^k$ ,  $\mathcal{P}_{\text{RotRot}^\top, \Gamma_t}^k$ ,  $\mathcal{P}_{\text{RotRot}^\top, \Gamma_t}^{k+1, k}$ , and  $\mathcal{P}_{\text{Div}, \Gamma_t}^k$  are right inverses of  $\text{symGrad}$ ,  $\text{RotRot}^\top$ , and  $\text{Div}$ , respectively.

**Theorem 3.25** (bounded regular decompositions from bounded regular potentials). For  $k \geq 0$ , the bounded regular decompositions

$$\begin{aligned} \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega) &= \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega) + \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega) = \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega) + \text{RotRot}^\top \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) \\ &= R(\tilde{\mathcal{Q}}_{\text{Div}, \Gamma_t}^{k,1}) \dot{+} \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega) \\ &= R(\tilde{\mathcal{Q}}_{\text{Div}, \Gamma_t}^{k,1}) \dot{+} R(\tilde{\mathcal{N}}_{\text{Div}, \Gamma_t}^k), \\ \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega) &= \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) + \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega) = \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) + \text{symGrad} \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega) \\ &= R(\tilde{\mathcal{Q}}_{\text{RotRot}^\top, \Gamma_t}^{k,1}) \dot{+} \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega) \\ &= R(\tilde{\mathcal{Q}}_{\text{RotRot}^\top, \Gamma_t}^{k,1}) \dot{+} R(\tilde{\mathcal{N}}_{\text{RotRot}^\top, \Gamma_t}^k), \\ \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega) &= \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) + \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^{k+1}(\text{RotRot}^\top, \Omega) = \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) + \text{symGrad} \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) \\ &= R(\tilde{\mathcal{Q}}_{\text{RotRot}^\top, \Gamma_t}^{k+1, k, 1}) \dot{+} \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^{k+1}(\text{RotRot}^\top, \Omega) \\ &= R(\tilde{\mathcal{Q}}_{\text{RotRot}^\top, \Gamma_t}^{k+1, k, 1}) \dot{+} R(\tilde{\mathcal{N}}_{\text{RotRot}^\top, \Gamma_t}^{k+1, k}). \end{aligned}$$

hold with bounded linear regular decomposition operators

$$\begin{aligned} \tilde{\mathcal{Q}}_{\text{Div}, \Gamma_t}^{k,1} &:= \mathcal{P}_{\text{Div}, \Gamma_t}^k \text{Div}_{\mathbb{S}, \Gamma_t}^k : \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega) \rightarrow \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1}(\Omega), \\ \tilde{\mathcal{Q}}_{\text{RotRot}^\top, \Gamma_t}^{k,1} &:= \mathcal{P}_{\text{RotRot}^\top, \Gamma_t}^k \text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k} : \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega), \\ \tilde{\mathcal{Q}}_{\text{RotRot}^\top, \Gamma_t}^{k+1, k, 1} &:= \mathcal{P}_{\text{RotRot}^\top, \Gamma_t}^{k+1, k} \text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k} : \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega), \\ \tilde{\mathcal{N}}_{\text{Div}, \Gamma_t}^k &: \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega) \rightarrow \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{Div}, \Omega), \\ \tilde{\mathcal{N}}_{\text{RotRot}^\top, \Gamma_t}^k &: \mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega), \\ \tilde{\mathcal{N}}_{\text{RotRot}^\top, \Gamma_t}^{k+1, k} &: \mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\mathbb{S}, \Gamma_t, 0}^{k+1}(\text{RotRot}^\top, \Omega) \end{aligned}$$

satisfying

$$\begin{aligned} \text{id}_{\mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{Div}, \Omega)} &= \tilde{\mathcal{Q}}_{\text{Div}, \Gamma_t}^{k,1} + \tilde{\mathcal{N}}_{\text{Div}, \Gamma_t}^k, \\ \text{id}_{\mathbf{H}_{\mathbb{S}, \Gamma_t}^k(\text{RotRot}^\top, \Omega)} &= \tilde{\mathcal{Q}}_{\text{RotRot}^\top, \Gamma_t}^{k,1} + \tilde{\mathcal{N}}_{\text{RotRot}^\top, \Gamma_t}^k, \\ \text{id}_{\mathbf{H}_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega)} &= \tilde{\mathcal{Q}}_{\text{RotRot}^\top, \Gamma_t}^{k+1, k, 1} + \tilde{\mathcal{N}}_{\text{RotRot}^\top, \Gamma_t}^{k+1, k}. \end{aligned}$$



**Corollary 3.26** (bounded regular kernel decompositions). *For  $k \geq 0$ , the bounded regular kernel decompositions*

$$\begin{aligned} H_{\mathbb{S},\Gamma_t,0}^k(\text{Div}, \Omega) &= H_{\mathbb{S},\Gamma_t,0}^{k+1}(\text{Div}, \Omega) + \text{RotRot}^\top H_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), \\ H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega) &= H_{\mathbb{S},\Gamma_t,0}^{k+2}(\text{RotRot}^\top, \Omega) + \text{symGrad} H_{\Gamma_t}^{k+1}(\Omega) \end{aligned}$$

hold.

*Remark 3.27* (bounded regular decompositions from bounded regular potentials). It holds

$$\text{Div} \tilde{Q}_{\text{Div}\mathbb{S},\Gamma_t}^{k,1} = \text{Div} Q_{\text{Div}\mathbb{S},\Gamma_t}^{k,1} = \text{Div}_{\mathbb{S},\Gamma_t}^k$$

and hence  $H_{\mathbb{S},\Gamma_t,0}^k(\text{Div}, \Omega)$  is invariant under  $Q_{\text{Div}\mathbb{S},\Gamma_t}^{k,1}$  and  $\tilde{Q}_{\text{Div}\mathbb{S},\Gamma_t}^{k,1}$ . Analogously,

$$\begin{aligned} \text{RotRot}^\top \tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1} &= \text{RotRot}^\top Q_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1} = \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}, \\ \text{RotRot}^\top \tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k,1} &= \text{RotRot}^\top Q_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k,1} = \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k}. \end{aligned}$$

Thus,  $H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega)$  and  $H_{\mathbb{S},\Gamma_t,0}^{k+1}(\text{RotRot}^\top, \Omega)$  are invariant under  $Q_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1}$  and  $Q_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k,1}$ , respectively. Moreover,

$$R(\tilde{Q}_{\text{Div}\mathbb{S},\Gamma_t}^{k,1}) = R(\mathcal{P}_{\text{Div}\mathbb{S},\Gamma_t}^k), \tilde{Q}_{\text{Div}\mathbb{S},\Gamma_t}^{k,1} = Q_{\text{Div}\mathbb{S},\Gamma_t}^{k,1} (\text{Div}_{\mathbb{S},\Gamma_t}^k)_\perp^{-1} \text{Div}_{\mathbb{S},\Gamma_t}^k.$$

Therefore, we have  $\tilde{Q}_{\text{Div}\mathbb{S},\Gamma_t}^{k,1} |_{D((\text{Div}_{\mathbb{S},\Gamma_t}^k)_\perp)} = Q_{\text{Div}\mathbb{S},\Gamma_t}^{k,1} |_{D((\text{Div}_{\mathbb{S},\Gamma_t}^k)_\perp)}$  and thus  $\tilde{Q}_{\text{Div}\mathbb{S},\Gamma_t}^{k,1}$  may differ from  $Q_{\text{Div}\mathbb{S},\Gamma_t}^{k,1}$  only on  $N(\text{Div}_{\mathbb{S},\Gamma_t}^k) = H_{\mathbb{S},\Gamma_t,0}^k(\text{Div}, \Omega)$ . Analogously,

$$\begin{aligned} R(\tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1}) &= R(\mathcal{P}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^k), & \tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1} &= Q_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1} (\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k})_\perp^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}, \\ R(\tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k,1}) &= R(\mathcal{P}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k}), & \tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k,1} &= Q_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k,1} (\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k})_\perp^{-1} \text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k}. \end{aligned}$$

Hence,

$$\begin{aligned} \tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1} |_{D((\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k})_\perp)} &= Q_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1} |_{D((\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k})_\perp)}, \\ \tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k,1} |_{D((\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k})_\perp)} &= Q_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k,1} |_{D((\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k})_\perp)} \end{aligned}$$

and thus  $\tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1}$  and  $\tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k,1}$  may differ from  $Q_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1}$  and  $Q_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k+1,k,1}$  only on the kernels  $N(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}) = H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega)$  and  $N(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k}) = H_{\mathbb{S},\Gamma_t,0}^{k+1}(\text{RotRot}^\top, \Omega)$ , respectively.

*Remark 3.28* (projections). Recall Theorem 3.25, for example, for  $\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}$

$$H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) = R(\tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1}) \dot{+} R(\tilde{\mathcal{N}}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^k).$$

- (i)  $\tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1}, \tilde{\mathcal{N}}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^k = 1 - \tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1}$  are projections.
- (i')  $\tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1} \tilde{\mathcal{N}}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^k = \tilde{\mathcal{N}}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^k \tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1} = 0$ .
- (ii) For  $I_\pm := \tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1} \pm \tilde{\mathcal{N}}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^k$ , it holds  $I_+ = I_-^2 = \text{id}_{H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega)}$ . Therefore,  $I_+, I_-^2$ , as well as  $I_- = 2\tilde{Q}_{\text{RotRot}^\top\mathbb{S},\Gamma_t}^{k,1} - \text{id}_{H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega)}$  are topological isomorphisms on  $H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega)$ .

(iii) There exists  $c > 0$  such that for all  $S \in \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega)$

$$\begin{aligned} c|\tilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1} S|_{\mathbf{H}_{\mathbb{S}}^{k+2}(\Omega)} &\leq |\text{RotRot}^\top S|_{\mathbf{H}_{\mathbb{S}}^k(\Omega)} \leq |S|_{\mathbf{H}_{\mathbb{S}}^k(\text{RotRot}^\top, \Omega)}, \\ |\tilde{\mathcal{N}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^k S|_{\mathbf{H}_{\mathbb{S}}^k(\Omega)} &\leq |S|_{\mathbf{H}_{\mathbb{S}}^k(\Omega)} + |\tilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1} S|_{\mathbf{H}_{\mathbb{S}}^k(\Omega)}. \end{aligned}$$

(iii') For  $S \in \mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega)$ , we have  $\tilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1} S = 0$  and  $\tilde{\mathcal{N}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^k S = S$ . In particular,  $\tilde{\mathcal{N}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^k$  is onto.

Similar results to (i)–(iii') hold for  $\text{Div}_{\mathbb{S},\Gamma_t}^k$  and also  $\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k}$ . In particular,  $\tilde{\mathcal{Q}}_{\text{Div}_{\mathbb{S},\Gamma_t}^k}^{k,1}$ ,  $\tilde{\mathcal{N}}_{\text{Div}_{\mathbb{S},\Gamma_t}^k}^k$ , and  $\tilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k,1}$ ,  $\tilde{\mathcal{N}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k}$  are projections and there exists  $c > 0$  such that for all  $T \in \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\text{Div}, \Omega)$  and all  $S \in \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1,k}(\text{RotRot}^\top, \Omega)$

$$|\tilde{\mathcal{Q}}_{\text{Div}_{\mathbb{S},\Gamma_t}^k}^{k,1} T|_{\mathbf{H}_{\mathbb{S}}^{k+1}(\Omega)} \leq c|\text{Div} T|_{\mathbf{H}^k(\Omega)}, \quad |\tilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k,1} S|_{\mathbf{H}_{\mathbb{S}}^{k+2}(\Omega)} \leq c|\text{RotRot}^\top S|_{\mathbf{H}_{\mathbb{S}}^k(\Omega)}.$$

Corollary 3.26 shows the following:

**Corollary 3.29** (bounded regular higher order kernel decompositions). *For  $k, \ell \geq 0$ , the bounded regular kernel decompositions*

$$\begin{aligned} N(\text{Div}_{\mathbb{S},\Gamma_t}^k) &= \mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{Div}, \Omega) = \mathbf{H}_{\mathbb{S},\Gamma_t,0}^\ell(\text{Div}, \Omega) + \text{RotRot}^\top \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+2}(\Omega), \\ N(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}) &= \mathbf{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega) = \mathbf{H}_{\mathbb{S},\Gamma_t,0}^\ell(\text{RotRot}^\top, \Omega) + \text{symGrad} \mathbf{H}_{\mathbb{S},\Gamma_t}^{k+1}(\Omega) \end{aligned}$$

hold. In particular, for  $k = 0$  and all  $\ell \geq 0$ ,

$$\begin{aligned} N(\text{Div}_{\mathbb{S},\Gamma_t}) &= \mathbf{H}_{\mathbb{S},\Gamma_t,0}(\text{Div}, \Omega) = \mathbf{H}_{\mathbb{S},\Gamma_t,0}^\ell(\text{Div}, \Omega) + \text{RotRot}^\top \mathbf{H}_{\mathbb{S},\Gamma_t}^2(\Omega), \\ N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top) &= \mathbf{H}_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^\top, \Omega) = \mathbf{H}_{\mathbb{S},\Gamma_t,0}^\ell(\text{RotRot}^\top, \Omega) + \text{symGrad} \mathbf{H}_{\mathbb{S},\Gamma_t}^1(\Omega). \end{aligned}$$

### 3.4 | Dirichlet/Neumann fields

From Theorem 3.15 (iv), we recall the orthonormal Helmholtz-type decompositions (for  $\mu = 1$ )

$$\begin{aligned} \mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega) &= R(\text{symGrad}_{\Gamma_t}) \oplus_{\mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega)} N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon) \\ &= N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top) \oplus_{\mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega)} R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top) \\ &= R(\text{symGrad}_{\Gamma_t}) \oplus_{\mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega)} \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) \oplus_{\mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega)} R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top), \\ N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top) &= R(\text{symGrad}_{\Gamma_t}) \oplus_{\mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega)} \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega), \\ N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon) &= \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) \oplus_{\mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega)} R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top). \end{aligned} \tag{13}$$

Let us denote the  $\mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega)$ -orthonormal projector onto  $N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon)$  and  $N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top)$  by

$$\pi_{\text{Div}} : \mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega) \rightarrow N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon), \quad \pi_{\text{RotRot}^\top} : \mathbf{L}_{\mathbb{S},\varepsilon}^2(\Omega) \rightarrow N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top),$$

respectively. Then

$$\begin{aligned} \pi_{\text{Div}}|_{N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top)} &: N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top) \rightarrow \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega), \\ \pi_{\text{RotRot}^\top}|_{N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon)} &: N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon) \rightarrow \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) \end{aligned}$$

are onto. Moreover,

$$\begin{aligned} \pi_{\text{Div}}|_{R(\text{symGrad}_{\Gamma_t})} &= 0, & \pi_{\text{RotRot}^\top}|_{R(\varepsilon^{-1}\text{RotRot}_{\mathbb{S},\Gamma_n}^\top)} &= 0, \\ \pi_{\text{Div}}|_{\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega)} &= \text{id}_{\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega)}, & \pi_{\text{RotRot}^\top}|_{\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega)} &= \text{id}_{\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega)}. \end{aligned}$$

Therefore, by Corollary 3.29 and for all  $\ell \geq 0$

$$\begin{aligned} \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) &= \pi_{\text{Div}}N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top) = \pi_{\text{Div}}\mathcal{H}_{\mathbb{S},\Gamma_t,0}^\ell(\text{RotRot}^\top, \Omega), \\ \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) &= \pi_{\text{RotRot}^\top}N(\text{Div}_{\mathbb{S},\Gamma_n}\varepsilon) = \pi_{\text{RotRot}^\top}\varepsilon^{-1}\mathcal{H}_{\mathbb{S},\Gamma_n,0}^\ell(\text{Div}, \Omega), \end{aligned}$$

where we have used  $N(\text{Div}_{\mathbb{S},\Gamma_n}\varepsilon) = \varepsilon^{-1}\mathcal{H}_{\mathbb{S},\Gamma_n,0}(\text{Div}, \Omega)$ . Hence, with

$$\mathcal{H}_{\mathbb{S},\Gamma_t,0}^\infty(\text{RotRot}^\top, \Omega) := \bigcap_{k \geq 0} \mathcal{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega), \quad \mathcal{H}_{\mathbb{S},\Gamma_n,0}^\infty(\text{Div}, \Omega) := \bigcap_{k \geq 0} \mathcal{H}_{\mathbb{S},\Gamma_n,0}^k(\text{Div}, \Omega),$$

we have the following result:

**Theorem 3.30** (smooth pre-bases of Dirichlet/Neumann fields). *Let  $d_{\Omega,\Gamma_t} := \dim \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega)$ . Then*

$$\pi_{\text{Div}}\mathcal{H}_{\mathbb{S},\Gamma_t,0}^\infty(\text{RotRot}^\top, \Omega) = \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) = \pi_{\text{RotRot}^\top}\varepsilon^{-1}\mathcal{H}_{\mathbb{S},\Gamma_n,0}^\infty(\text{Div}, \Omega).$$

Moreover, there exists a smooth  $\text{RotRot}^\top$ -pre-basis and a smooth  $\text{Div}$ -pre-basis of  $\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega)$ ; that is, there are linear independent smooth fields

$$\begin{aligned} \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega) &:= \{B_{\mathbb{S},\Gamma_t,\ell}^{\text{RotRot}^\top}\}_{\ell=1}^{d_{\Omega,\Gamma_t}} \subset \mathcal{H}_{\mathbb{S},\Gamma_t,0}^\infty(\text{RotRot}^\top, \Omega), \\ \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega) &:= \{B_{\mathbb{S},\Gamma_n,\ell}^{\text{Div}}\}_{\ell=1}^{d_{\Omega,\Gamma_t}} \subset \mathcal{H}_{\mathbb{S},\Gamma_n,0}^\infty(\text{Div}, \Omega), \end{aligned}$$

such that  $\pi_{\text{Div}}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega)$  and  $\pi_{\text{RotRot}^\top}\varepsilon^{-1}\mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega)$  are both bases of  $\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega)$ . In particular,

$$\text{Lin}\pi_{\text{Div}}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega) = \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) = \text{Lin}\pi_{\text{RotRot}^\top}\varepsilon^{-1}\mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega).$$

Note that  $(1 - \pi_{\text{Div}})$  and  $(1 - \pi_{\text{RotRot}^\top})$  are the  $L^2_{\mathbb{S},\varepsilon}(\Omega)$ -orthonormal projectors onto the ranges  $R(\text{symGrad}_{\Gamma_t})$  and  $R(\varepsilon^{-1}\text{RotRot}_{\mathbb{S},\Gamma_n}^\top)$ , respectively, that is,

$$(1 - \pi_{\text{Div}}) : L^2_{\mathbb{S},\varepsilon}(\Omega) \rightarrow R(\text{symGrad}_{\Gamma_t}), \quad (1 - \pi_{\text{RotRot}^\top}) : L^2_{\mathbb{S},\varepsilon}(\Omega) \rightarrow R(\varepsilon^{-1}\text{RotRot}_{\mathbb{S},\Gamma_n}^\top).$$

By (13) and Theorems 3.16 and 3.30, we have, for example,

$$\begin{aligned} \mathcal{H}_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^\top, \Omega) &= R(\text{symGrad}_{\Gamma_t}) \oplus_{L^2_{\mathbb{S},\varepsilon}(\Omega)} \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) \\ &= R(\text{symGrad}_{\Gamma_t}) \oplus_{L^2_{\mathbb{S},\varepsilon}(\Omega)} \text{Lin}\pi_{\text{Div}}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega) \\ &= R(\text{symGrad}_{\Gamma_t}) + (\pi_{\text{Div}} - 1)\text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega) + \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega) \\ &= R(\text{symGrad}_{\Gamma_t}) + \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega), \\ \mathcal{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega) &= R(\text{symGrad}_{\Gamma_t}) \cap \mathcal{H}_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega) + \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega), \\ &= R(\text{symGrad}_{\Gamma_t}^k) + \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega). \end{aligned} \tag{14}$$

Similarly, we obtain a decomposition of  $H_{\mathbb{S},\Gamma_n,0}^k(\text{Div}, \Omega)$  using  $\mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega)$ . We conclude the following:

**Theorem 3.31** (bounded regular direct decompositions). *Let  $k \geq 0$ . Then the bounded regular direct decompositions*

$$\begin{aligned} H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) &= R(\tilde{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1}) \dot{+} H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega), \\ H_{\mathbb{S},\Gamma_t}^{k+1,k}(\text{RotRot}^\top, \Omega) &= R(\tilde{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k,1}) \dot{+} H_{\mathbb{S},\Gamma_t,0}^{k+1}(\text{RotRot}^\top, \Omega), \\ H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega) &= \text{symGrad } H_{\Gamma_t}^{k+1}(\Omega) \dot{+} \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega), \\ H_{\mathbb{S},\Gamma_n}^k(\text{Div}, \Omega) &= R(\tilde{Q}_{\text{Div}_{\mathbb{S},\Gamma_n}}^{k,1}) \dot{+} H_{\mathbb{S},\Gamma_n,0}^k(\text{Div}, \Omega), \\ H_{\mathbb{S},\Gamma_n,0}^k(\text{Div}, \Omega) &= \text{RotRot}^\top H_{\mathbb{S},\Gamma_n}^{k+2}(\Omega) \dot{+} \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega) \end{aligned}$$

hold. Note that  $R(\tilde{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1}), R(\tilde{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k,1}) \subset H_{\mathbb{S},\Gamma_t}^{k+2}(\Omega)$  and  $R(\tilde{Q}_{\text{Div}_{\mathbb{S},\Gamma_n}}^{k,1}) \subset H_{\mathbb{S},\Gamma_n}^{k+1}(\Omega)$ .

*Remark 3.32.* (bounded regular direct decompositions) In particular, for  $k = 0$ ,

$$\begin{aligned} H_{\mathbb{S},\Gamma_t}(\text{RotRot}^\top, \Omega) &= R(\tilde{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{0,1}) \dot{+} H_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^\top, \Omega), \\ H_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^\top, \Omega) &= \text{symGrad } H_{\Gamma_t}^1(\Omega) \dot{+} \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega) \\ &= \text{symGrad } H_{\Gamma_t}^1(\Omega) \oplus_{L_{\mathbb{S},\epsilon}^2(\Omega)} \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\epsilon}(\Omega), \\ H_{\mathbb{S},\Gamma_n}(\text{Div}, \Omega) &= R(\tilde{Q}_{\text{Div}_{\mathbb{S},\Gamma_n}}^{0,1}) \dot{+} H_{\mathbb{S},\Gamma_n,0}(\text{Div}, \Omega), \\ \epsilon^{-1} H_{\mathbb{S},\Gamma_n,0}(\text{Div}, \Omega) &= \epsilon^{-1} \text{RotRot}^\top H_{\mathbb{S},\Gamma_n}^2(\Omega) \dot{+} \epsilon^{-1} \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega) \\ &= \epsilon^{-1} \text{RotRot}^\top H_{\mathbb{S},\Gamma_n}^2(\Omega) \oplus_{L_{\mathbb{S},\epsilon}^2(\Omega)} \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\epsilon}(\Omega) \end{aligned}$$

and

$$\begin{aligned} L_{\mathbb{S},\epsilon}^2(\Omega) &= H_{\mathbb{S},\Gamma_t,0}(\text{RotRot}^\top, \Omega) \oplus_{L_{\mathbb{S},\epsilon}^2(\Omega)} \epsilon^{-1} \text{RotRot}^\top H_{\mathbb{S},\Gamma_n}^2(\Omega) \\ &= \text{symGrad } H_{\Gamma_t}^1(\Omega) \oplus_{L_{\mathbb{S},\epsilon}^2(\Omega)} \epsilon^{-1} H_{\mathbb{S},\Gamma_n,0}(\text{Div}, \Omega). \end{aligned}$$

*Proof of Theorem 3.31.* Theorem 3.25 and (14) show

$$\begin{aligned} H_{\mathbb{S},\Gamma_t}^k(\text{RotRot}^\top, \Omega) &= R(\tilde{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k,1}) \dot{+} H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega), \\ H_{\mathbb{S},\Gamma_t}^{k+1,k}(\text{RotRot}^\top, \Omega) &= R(\tilde{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k,1}) \dot{+} H_{\mathbb{S},\Gamma_t,0}^{k+1}(\text{RotRot}^\top, \Omega), \\ H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega) &= \text{symGrad } H_{\Gamma_t}^{k+1}(\Omega) + \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega). \end{aligned}$$

To prove the directness of the third sum, let

$$\sum_{\ell=1}^{d_{\Omega,\Gamma_t}} \lambda_\ell \mathcal{B}_{\mathbb{S},\Gamma_t,\ell}^{\text{RotRot}^\top} \in \text{symGrad } H_{\Gamma_t}^{k+1}(\Omega) \cap \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega).$$

Then  $0 = \sum_{\ell} \lambda_\ell \pi_{\text{Div}} \mathcal{B}_{\mathbb{S},\Gamma_t,\ell}^{\text{RotRot}^\top} \in \text{Lin}\pi_{\text{Div}} \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega)$  and hence  $\lambda_\ell = 0$  for all  $\ell$  as  $\pi_{\text{Div}} \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega)$  is a basis of  $\mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\epsilon}(\Omega)$  by Theorem 3.30. Concerning the boundedness of the decompositions, let

$$H_{\mathbb{S},\Gamma_t,0}^k(\text{RotRot}^\top, \Omega) \ni S = \text{symGrad } v + B, \quad v \in H_{\Gamma_t}^{k+1}(\Omega), \quad B \in \text{Lin}\mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega).$$

By Theorem 3.24  $\text{symGrad } v \in R(\text{symGrad}_{\Gamma_t}^k)$  and  $u := \mathcal{P}_{\text{symGrad}, \Gamma_t}^k \text{symGrad } v \in \mathbf{H}_{\Gamma_t}^{k+1}(\Omega)$  solves  $\text{symGrad } u = \text{symGrad } v$  with  $|u|_{\mathbf{H}^{k+1}(\Omega)} \leq c|\text{symGrad } v|_{\mathbf{H}_S^k(\Omega)}$ . Therefore,

$$|u|_{\mathbf{H}^{k+1}(\Omega)} + |B|_{\mathbf{H}_S^k(\Omega)} \leq c \left( |\text{symGrad } v|_{\mathbf{H}_S^k(\Omega)} + |B|_{\mathbf{H}_S^k(\Omega)} \right) \leq c \left( |S|_{\mathbf{H}_S^k(\Omega)} + |B|_{\mathbf{H}_S^k(\Omega)} \right).$$

Note that the mapping

$$I_{\pi, \text{Div}} : \text{Lin } \mathcal{B}_{S, \Gamma_t}^{\text{RotRot}^\top}(\Omega) \rightarrow \text{Lin } \pi_{\text{Div}} \mathcal{B}_{S, \Gamma_t}^{\text{RotRot}^\top}(\Omega) = \mathcal{H}_{S, \Gamma_t, \Gamma_n, \epsilon}(\Omega); \mathcal{B}_{S, \Gamma_t, \ell}^{\text{RotRot}^\top} \mapsto \pi_{\text{Div}} \mathcal{B}_{S, \Gamma_t, \ell}^{\text{RotRot}^\top}$$

is a topological isomorphism (between finite dimensional spaces and with arbitrary norms). Thus,

$$|B|_{\mathbf{H}_S^k(\Omega)} \leq c|B|_{L_S^2(\Omega)} \leq c|\pi_{\text{Div}} B|_{L_S^2(\Omega)} = c|\pi_{\text{Div}} S|_{L_S^2(\Omega)} \leq c|S|_{L_S^2(\Omega)} \leq c|S|_{\mathbf{H}_S^k(\Omega)}.$$

Finally, we see  $S = \text{symGrad } u + B \in \text{symGrad } \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \dot{+} \text{Lin } \mathcal{B}_{S, \Gamma_t}^{\text{RotRot}^\top}(\Omega)$  and

$$|u|_{\mathbf{H}^{k+1}(\Omega)} + |B|_{\mathbf{H}_S^k(\Omega)} \leq c|S|_{\mathbf{H}_S^k(\Omega)}.$$

The other assertions for Div follow analogously. □

*Remark 3.33* (bounded regular direct decompositions). By Theorem 3.31, we have, for example,

$$\begin{aligned} \mathbf{H}_{S, \Gamma_t}^k(\text{RotRot}^\top, \Omega) &= R(\tilde{\mathcal{Q}}_{\text{RotRot}_{S, \Gamma_t}^\top}^{k,1}) \dot{+} \text{Lin } \mathcal{B}_{S, \Gamma_t}^{\text{RotRot}^\top}(\Omega) \dot{+} \text{symGrad } \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \\ &= \mathbf{H}_{S, \Gamma_t}^{k+2}(\Omega) + \text{symGrad } \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \end{aligned}$$

with bounded linear regular direct decomposition operators

$$\begin{aligned} \hat{\mathcal{Q}}_{\text{RotRot}_{S, \Gamma_t}^\top}^{k,1} &: \mathbf{H}_{S, \Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow R(\tilde{\mathcal{Q}}_{\text{RotRot}_{S, \Gamma_t}^\top}^{k,1}) \subset \mathbf{H}_{S, \Gamma_t}^{k+2}(\Omega), \\ \hat{\mathcal{Q}}_{\text{RotRot}_{S, \Gamma_t}^\top}^{k,\infty} &: \mathbf{H}_{S, \Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow \text{Lin } \mathcal{B}_{S, \Gamma_t}^{\text{RotRot}^\top}(\Omega) \subset \mathbf{H}_{S, \Gamma_t, 0}^\infty(\text{RotRot}^\top, \Omega) \subset \mathbf{H}_{S, \Gamma_t}^{k+2}(\Omega), \\ \hat{\mathcal{Q}}_{\text{RotRot}_{S, \Gamma_t}^\top}^{k,0} &: \mathbf{H}_{S, \Gamma_t}^k(\text{RotRot}^\top, \Omega) \rightarrow \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \end{aligned}$$

satisfying  $\hat{\mathcal{Q}}_{\text{RotRot}_{S, \Gamma_t}^\top}^{k,1} + \hat{\mathcal{Q}}_{\text{RotRot}_{S, \Gamma_t}^\top}^{k,\infty} + \text{symGrad } \hat{\mathcal{Q}}_{\text{RotRot}_{S, \Gamma_t}^\top}^{k,0} = \text{id}_{\mathbf{H}_{S, \Gamma_t}^k(\text{RotRot}^\top, \Omega)}$ .

A closer inspection of the latter proof allows for a more precise description of these bounded decomposition operators. For this, let  $S \in \mathbf{H}_{S, \Gamma_t}^k(\text{RotRot}^\top, \Omega)$ . According to Theorem 3.25 and Remark 3.28, we decompose

$$S = S_R + S_N := \tilde{\mathcal{Q}}_{\text{RotRot}_{S, \Gamma_t}^\top}^{k,1} S + \tilde{\mathcal{N}}_{\text{RotRot}_{S, \Gamma_t}^\top}^k S \in R(\tilde{\mathcal{Q}}_{\text{RotRot}_{S, \Gamma_t}^\top}^{k,1}) \dot{+} R(\tilde{\mathcal{N}}_{\text{RotRot}_{S, \Gamma_t}^\top}^k)$$

with  $R(\tilde{\mathcal{N}}_{\text{RotRot}_{S, \Gamma_t}^\top}^k) = \mathbf{H}_{S, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega) = N(\text{RotRot}_{S, \Gamma_t}^{\top, k})$ . By Theorem 3.31, we further decompose

$$\mathbf{H}_{S, \Gamma_t, 0}^k(\text{RotRot}^\top, \Omega) \ni S_N = \text{symGrad } u + B \in \text{symGrad } \mathbf{H}_{\Gamma_t}^{k+1}(\Omega) \dot{+} \text{Lin } \mathcal{B}_{S, \Gamma_t}^{\text{RotRot}^\top}(\Omega).$$

Then  $\pi_{\text{Div}} S_N = \pi_{\text{Div}} B \in \mathcal{H}_{\mathbb{S}, \Gamma_t, \Gamma_n, \epsilon}(\Omega)$  and thus  $B = I_{\pi, \text{Div}}^{-1} \pi_{\text{Div}} S_N \in \text{Lin} \mathcal{B}_{\mathbb{S}, \Gamma_t}^{\text{RotRot}^\top}(\Omega)$ . Therefore,  $u = \mathcal{P}_{\text{symGrad}, \Gamma_t}^k \text{symGrad } u = \mathcal{P}_{\text{symGrad}, \Gamma_t}^k (S_N - B) = \mathcal{P}_{\text{symGrad}, \Gamma_t}^k (1 - I_{\pi, \text{Div}}^{-1} \pi_{\text{Div}}) S_N$ . Finally, we see

$$\begin{aligned} \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k,1} &= \widetilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k,1} = \mathcal{P}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^k \text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k} = \mathcal{Q}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k,1} (\text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k})^{-1} \text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k}, \\ \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k,\infty} &= I_{\pi, \text{Div}}^{-1} \pi_{\text{Div}} \widetilde{\mathcal{N}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^k, \\ \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k,0} &= \mathcal{P}_{\text{symGrad}, \Gamma_t}^k (1 - I_{\pi, \text{Div}}^{-1} \pi_{\text{Div}}) \widetilde{\mathcal{N}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^k \end{aligned}$$

with  $\widetilde{\mathcal{N}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^k = 1 - \widetilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k,1}$ . Analogously, we have

$$\begin{aligned} \mathbb{H}_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega) &= R(\widetilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, 1}) \dot{+} \text{Lin} \mathcal{B}_{\mathbb{S}, \Gamma_t}^{\text{RotRot}^\top}(\Omega) \dot{+} \text{symGrad } \mathbb{H}_{\Gamma_t}^{k+2}(\Omega) \\ &= \mathbb{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega) + \text{symGrad } \mathbb{H}_{\Gamma_t}^{k+2}(\Omega), \\ \mathbb{H}_{\mathbb{S}, \Gamma_n}^k(\text{Div}, \Omega) &= R(\widetilde{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,1}) \dot{+} \text{Lin} \mathcal{B}_{\mathbb{S}, \Gamma_n}^{\text{Div}}(\Omega) \dot{+} \text{RotRot}^\top \mathbb{H}_{\mathbb{S}, \Gamma_n}^{k+2}(\Omega) \\ &= \mathbb{H}_{\mathbb{S}, \Gamma_n}^{k+1}(\Omega) + \text{RotRot}^\top \mathbb{H}_{\mathbb{S}, \Gamma_n}^{k+2}(\Omega) \end{aligned}$$

with bounded linear regular direct decomposition operators

$$\begin{aligned} \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, 1} &: \mathbb{H}_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega) \rightarrow R(\widetilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, 1}) \subset \mathbb{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega), \\ \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, \infty} &: \mathbb{H}_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega) \rightarrow \text{Lin} \mathcal{B}_{\mathbb{S}, \Gamma_t}^{\text{RotRot}^\top}(\Omega) \subset \mathbb{H}_{\mathbb{S}, \Gamma_t, 0}^\infty(\text{RotRot}^\top, \Omega) \subset \mathbb{H}_{\mathbb{S}, \Gamma_t}^{k+2}(\Omega), \\ \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, 0} &: \mathbb{H}_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega) \rightarrow \mathbb{H}_{\Gamma_t}^{k+2}(\Omega), \\ \widehat{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,1} &: \mathbb{H}_{\mathbb{S}, \Gamma_n}^k(\text{Div}, \Omega) \rightarrow R(\widetilde{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,1}) \subset \mathbb{H}_{\mathbb{S}, \Gamma_n}^{k+1}(\Omega), \\ \widehat{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,\infty} &: \mathbb{H}_{\mathbb{S}, \Gamma_n}^k(\text{Div}, \Omega) \rightarrow \text{Lin} \mathcal{B}_{\mathbb{S}, \Gamma_n}^{\text{Div}}(\Omega) \subset \mathbb{H}_{\mathbb{S}, \Gamma_n, 0}^\infty(\text{Div}, \Omega) \subset \mathbb{H}_{\mathbb{S}, \Gamma_n}^{k+1}(\Omega), \\ \widehat{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,0} &: \mathbb{H}_{\mathbb{S}, \Gamma_n}^k(\text{Div}, \Omega) \rightarrow \mathbb{H}_{\mathbb{S}, \Gamma_n}^{k+2}(\Omega) \end{aligned}$$

satisfying

$$\begin{aligned} \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, 1} + \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, \infty} + \text{symGrad} \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, 0} &= \text{id}_{\mathbb{H}_{\mathbb{S}, \Gamma_t}^{k+1, k}(\text{RotRot}^\top, \Omega)}, \\ \widehat{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,1} + \widehat{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,\infty} + \text{RotRot}^\top \widehat{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,0} &= \text{id}_{\mathbb{H}_{\mathbb{S}, \Gamma_n}^k(\text{Div}, \Omega)} \end{aligned}$$

and

$$\begin{aligned} \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, 1} &= \widetilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, 1} = \mathcal{P}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k} \text{RotRot}_{\mathbb{S}, \Gamma_t}^{\top, k+1, k}, \\ \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, \infty} &= I_{\pi, \text{Div}}^{-1} \pi_{\text{Div}} \widetilde{\mathcal{N}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k}, \\ \widehat{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k, 0} &= \mathcal{P}_{\text{symGrad}, \Gamma_t}^{k+1} (1 - I_{\pi, \text{Div}}^{-1} \pi_{\text{Div}}) \widetilde{\mathcal{N}}_{\text{RotRot}_{\mathbb{S}, \Gamma_t}^\top}^{k+1, k}, \\ \widehat{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,1} &= \widetilde{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,1} = \mathcal{P}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^k \text{Div}_{\mathbb{S}, \Gamma_n}^k, \\ \widehat{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,\infty} &= I_{\pi, \text{RotRot}^\top}^{-1} \pi_{\text{RotRot}^\top} \widetilde{\mathcal{N}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^k, \\ \widehat{\mathcal{Q}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^{k,0} &= \mathcal{P}_{\text{RotRot}_{\mathbb{S}, \Gamma_n}^\top}^k (1 - I_{\pi, \text{RotRot}^\top}^{-1} \pi_{\text{RotRot}^\top}) \widetilde{\mathcal{N}}_{\text{Div}_{\mathbb{S}, \Gamma_n}}^k \end{aligned}$$

with

$$\begin{aligned} \widetilde{\mathcal{N}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k} &= 1 - \widetilde{\mathcal{Q}}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k,1}, & \widetilde{\mathcal{N}}_{\text{Div}_{\mathbb{S},\Gamma_n}}^k &= 1 - \widehat{\mathcal{Q}}_{\text{Div}_{\mathbb{S},\Gamma_n}}^{k,1}, \\ \mathcal{P}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k} &= \mathcal{Q}_{\text{RotRot}_{\mathbb{S},\Gamma_t}^\top}^{k+1,k,1} (\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k+1,k})_{\perp}^{-1}, & \mathcal{P}_{\text{Div}_{\mathbb{S},\Gamma_n}}^k &= \mathcal{Q}_{\text{Div}_{\mathbb{S},\Gamma_n}}^{k,1} (\text{Div}_{\mathbb{S},\Gamma_n}^k)_{\perp}^{-1}, \end{aligned}$$

and

$$I_{\pi, \text{RotRot}^\top} : \text{Lin} \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega) \rightarrow \text{Lin} \pi_{\text{RotRot}^\top} \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega) = \mathcal{H}_{\mathbb{S},\Gamma_n,\Gamma_n,\varepsilon}(\Omega); \mathcal{B}_{\mathbb{S},\Gamma_n,\ell}^{\text{Div}} \mapsto \pi_{\text{RotRot}^\top} \mathcal{B}_{\mathbb{S},\Gamma_n,\ell}^{\text{Div}}.$$

Noting

$$R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top)_{\perp} \mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega) \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega), \quad R(\text{symGrad}_{\Gamma_t})_{\perp} \mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega) \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega), \tag{15}$$

we see the following:

**Theorem 3.34** (alternative Dirichlet/Neumann projections). *It holds*

$$\begin{aligned} \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) \cap \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega)_{\perp} \mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega) &= \{0\}, \\ N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon) \cap \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega)_{\perp} \mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega) &= R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top), \\ \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) \cap \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega)_{\perp} \mathcal{L}_{\mathbb{S}}^2(\Omega) &= \{0\}, \\ N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top) \cap \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega)_{\perp} \mathcal{L}_{\mathbb{S}}^2(\Omega) &= R(\text{symGrad}_{\Gamma_t}). \end{aligned}$$

Moreover, for all  $k \geq 0$ ,

$$\begin{aligned} N(\text{Div}_{\mathbb{S},\Gamma_n}^k \varepsilon) \cap \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega)_{\perp} \mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega) &= R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k}) = \varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top \mathcal{H}_{\mathbb{S},\Gamma_n}^{k+2}(\Omega), \\ N(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}) \cap \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega)_{\perp} \mathcal{L}_{\mathbb{S}}^2(\Omega) &= R(\text{symGrad}_{\Gamma_t}^k) = \text{symGrad}_{\Gamma_t} \mathcal{H}_{\Gamma_t}^{k+1}(\Omega). \end{aligned}$$

*Proof.* For  $k = 0$  and  $S \in \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) \cap \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^\top}(\Omega)_{\perp} \mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega)$ , we have

$$0 = \langle S, \mathcal{B}_{\mathbb{S},\Gamma_t,\ell}^{\text{RotRot}^\top} \rangle_{\mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega)} = \langle \pi_{\text{Div}} S, \mathcal{B}_{\mathbb{S},\Gamma_t,\ell}^{\text{RotRot}^\top} \rangle_{\mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega)} = \langle S, \pi_{\text{Div}} \mathcal{B}_{\mathbb{S},\Gamma_t,\ell}^{\text{RotRot}^\top} \rangle_{\mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega)}$$

and hence  $S = 0$  by Theorem 3.30. Analogously, we see for  $S \in \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) \cap \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega)_{\perp} \mathcal{L}_{\mathbb{S}}^2(\Omega)$

$$0 = \langle S, \mathcal{B}_{\mathbb{S},\Gamma_n,\ell}^{\text{Div}} \rangle_{\mathcal{L}_{\mathbb{S}}^2(\Omega)} = \langle \pi_{\text{RotRot}^\top} S, \varepsilon^{-1} \mathcal{B}_{\mathbb{S},\Gamma_n,\ell}^{\text{Div}} \rangle_{\mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega)} = \langle S, \pi_{\text{RotRot}^\top} \varepsilon^{-1} \mathcal{B}_{\mathbb{S},\Gamma_n,\ell}^{\text{Div}} \rangle_{\mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega)}$$

and thus  $S = 0$  again by Theorem 3.30. According to (13), we can decompose

$$\begin{aligned} N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon) &= R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^\top) \oplus_{\mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega)} \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega), \\ N(\text{RotRot}_{\mathbb{S},\Gamma_t}^\top) &= R(\text{symGrad}_{\Gamma_t}) \oplus_{\mathcal{L}_{\mathbb{S},\varepsilon}^2(\Omega)} \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega), \end{aligned}$$

which shows by (15) the other two assertions. Let  $k \geq 0$ . The case  $k = 0$  and Theorem 3.16 show

$$\begin{aligned} N(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k}) \cap \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega)^{\perp_{L^2_{\mathbb{S}}(\Omega)}} &= \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\Omega) \cap N(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top}) \cap \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega)^{\perp_{L^2_{\mathbb{S}}(\Omega)}} \\ &= \mathbf{H}_{\mathbb{S},\Gamma_t}^k(\Omega) \cap R(\text{symGrad}_{\Gamma_t}) \\ &= R(\text{symGrad}_{\Gamma_t}^k) = \text{symGrad}_{\Gamma_t}^{k+1}(\Omega). \end{aligned}$$

Analogously,

$$\begin{aligned} N(\text{Div}_{\mathbb{S},\Gamma_n}^k \varepsilon) \cap \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^{\top}}(\Omega)^{\perp_{L^2_{\mathbb{S},\varepsilon}(\Omega)}} &= \varepsilon^{-1} \mathbf{H}_{\mathbb{S},\Gamma_n}^k(\Omega) \cap N(\text{Div}_{\mathbb{S},\Gamma_n} \varepsilon) \cap \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^{\top}}(\Omega)^{\perp_{L^2_{\mathbb{S},\varepsilon}(\Omega)}} \\ &= \varepsilon^{-1} \mathbf{H}_{\mathbb{S},\Gamma_n}^k(\Omega) \cap R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^{\top}) \\ &= R(\varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k}) = \varepsilon^{-1} \text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k+2}(\Omega), \end{aligned}$$

completing the proof. □

Theorem 3.31 implies the following:

**Theorem 3.35** (cohomology groups). *It holds*

$$\frac{N(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k})}{R(\text{symGrad}_{\Gamma_t}^k)} \cong \text{Lin} \mathcal{B}_{\mathbb{S},\Gamma_t}^{\text{RotRot}^{\top}}(\Omega) \cong \mathcal{H}_{\mathbb{S},\Gamma_t,\Gamma_n,\varepsilon}(\Omega) \cong \text{Lin} \mathcal{B}_{\mathbb{S},\Gamma_n}^{\text{Div}}(\Omega) \cong \frac{N(\text{Div}_{\mathbb{S},\Gamma_n}^k)}{R(\text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k})}.$$

*In particular, the dimensions of the cohomology groups (Dirichlet/Neumann fields) are independent of  $k$  and  $\varepsilon$  and it holds*

$$d_{\Omega,\Gamma_t} = \dim(N(\text{RotRot}_{\mathbb{S},\Gamma_t}^{\top,k})/R(\text{symGrad}_{\Gamma_t}^k)) = \dim(N(\text{Div}_{\mathbb{S},\Gamma_n}^k)/R(\text{RotRot}_{\mathbb{S},\Gamma_n}^{\top,k})).$$

## ACKNOWLEDGEMENTS

We thank the anonymous referee for careful reading and helpful suggestions to improve the quality of the paper. There are no funders to report for this submission. Open Access funding enabled and organized by Projekt DEAL.

## CONFLICT OF INTEREST

This work does not have any conflicts of interest.

## ORCID

Dirk Pauly  <https://orcid.org/0000-0003-4155-7297>

## REFERENCES

1. Pauly D, Schomburg M. Hilbert complexes with mixed boundary conditions—Part 1: de Rham complex. arXiv, 2021. <https://arxiv.org/abs/2106.03448>
2. Pauly D, Schomburg M. Hilbert complexes with mixed boundary conditions—Part 1: de Rham complex. *Math Methods Appl Sci.* 2022;45(5):2465-2507.
3. Arnold DN, Hu K. Complexes from complexes. arXiv, 2020. <https://arxiv.org/abs/2005.12437v1>
4. Pauly D. On the Maxwell constants in 3D. *Math Methods Appl Sci.* 2017;40(2):435-447.
5. Pauly D. A global div-curl-lemma for mixed boundary conditions in weak Lipschitz domains and a corresponding generalized  $A_0^*$ - $A_1$ -lemma in Hilbert spaces. *Anal (Berlin).* 2019;39:33-58.
6. Pauly D. On the Maxwell and Friedrichs/Poincare constants in ND. *Math Z.* 2019;293(3):957-987.
7. Pauly D. Solution theory, variational formulations, and functional a posteriori error estimates for general first order systems with applications to electro-magneto-statics and more. *Numer Funct Anal Optim.* 2020;41(1):16-112.



8. Pauly D, Zulehner W. On closed and exact Grad-grad- and div-Div-complexes, corresponding compact embeddings for tensor rotations, and a related decomposition result for biharmonic problems in 3D. arXiv, 2016. <https://arxiv.org/abs/1609.05873>
9. Pauly D, Zulehner W. The divDiv-complex and applications to biharmonic equations. *Appl Anal*. 2020;99(9):1579-1630.
10. Pauly D, Zulehner W. The elasticity complex: compact embeddings and regular decompositions. arXiv, 2020. <https://arxiv.org/abs/2001.11007>
11. Pauly D, Zulehner W. The elasticity complex: compact embeddings and regular decompositions. *Appl Anal*. 2022.
12. Bauer S, Pauly D, Schomburg M. The Maxwell compactness property in bounded weak Lipschitz domains with mixed boundary conditions. *SIAM J Math Anal*. 2016;48(4):2912-2943.
13. Neff P, Pauly D, Witsch K-J. Poincaré meets Korn via Maxwell: extending Korn's first inequality to incompatible tensor fields. *J Differ Equ*. 2015;258(4):1267-1302.
14. Jochmann F. A compactness result for vector fields with divergence and curl in  $L^q(\Omega)$  involving mixed boundary conditions. *Appl Anal*. 1997;66:189-203.
15. Picard R. An elementary proof for a compact imbedding result in generalized electromagnetic theory. *Math Z*. 1984;187:151-164.
16. Picard R, Weck N, Witsch K-J. Time-harmonic Maxwell equations in the exterior of perfectly conducting, irregular obstacles. *Anal (Munich)*. 2001;21:231-263.
17. Weber C. A local compactness theorem for Maxwell's equations. *Math Methods Appl Sci*. 1980;2:12-25.
18. Weck N. Maxwell's boundary value problems on Riemannian manifolds with nonsmooth boundaries. *J Math Anal Appl*. 1974;46:410-437.
19. Witsch K-J. A remark on a compactness result in electromagnetic theory. *Math Methods Appl Sci*. 1993;16:123-129.
20. Bauer S, Pauly D, Schomburg M. Weck's selection theorem: the Maxwell compactness property for bounded weak Lipschitz domains with mixed boundary conditions in arbitrary dimensions. arXiv, 2018. <https://arxiv.org/abs/1809.01192>
21. Bauer S, Pauly D, Schomburg M. Weck's selection theorem: the Maxwell compactness property for bounded weak Lipschitz domains with mixed boundary conditions in arbitrary dimensions. *Maxwell's Equations: Analysis and Numerics (Radon Series on Computational and Applied Mathematics)*: De Gruyter; 2019.
22. Pauly D, Waurick M. The index of some mixed order Dirac-type operators and generalised Dirichlet–Neumann tensor fields. arXiv, 2020. <https://arxiv.org/abs/2005.07996>
23. Pauly D, Waurick M. The index of some mixed order Dirac-type operators and generalised Dirichlet–Neumann tensor fields. *Math Z*. 2022;301(2):1739-1819.

**How to cite this article:** Pauly D, Schomburg M. Hilbert complexes with mixed boundary conditions—Part 2: Elasticity complex. *Math Meth Appl Sci*. 2022;45(16):8971-9005. doi:10.1002/mma.8242

## APPENDIX A: ELEMENTARY FORMULAS

From previous works<sup>8–11</sup> and references,<sup>22,23</sup> we have the following collection of formulas related to the elasticity and the biharmonic complex.

**Lemma A.1.** (lemma 12.10<sup>22,23</sup>) Let  $u, v, w$ , and  $S$  belong to  $C^\infty(\mathbb{R}^3)$ .

- $(\text{spn } v)w = v \times w = -(\text{spn } w)v$  and  $(\text{spn } v)(\text{spn}^{-1}S) = -Sv$ , if  $\text{sym } S = 0$
- $\text{sym spn } v = 0$  and  $\text{dev}(u \text{ id}) = 0$
- $\text{tr Grad } v = \text{div } v$  and  $2 \text{skw Grad } v = \text{spn rot } v$
- $\text{Div}(u \text{ id}) = \text{grad } u$  and  $\text{Rot}(u \text{ id}) = -\text{spn grad } u$ , in particular,  $\text{rot Div}(u \text{ id}) = 0$  and  $\text{rot spn}^{-1} \text{Rot}(u \text{ id}) = 0$  and  $\text{sym Rot}(u \text{ id}) = 0$
- $\text{Div spn } v = -\text{rot } v$  and  $\text{Div skw } S = -\text{rot spn}^{-1} \text{skw } S$ , in particular,  $\text{div Div skw } S = 0$
- $\text{Rot spn } v = (\text{div } v) \text{ id} - (\text{Grad } v)^\top$  and  $\text{Rot skw } S = (\text{div spn}^{-1} \text{skw } S) \text{ id} - (\text{Grad spn}^{-1} \text{skw } S)^\top$
- $\text{dev Rot spn } v = -(\text{dev Grad } v)^\top$
- $-2 \text{Rot sym Grad } v = 2 \text{Rot skw Grad } v = -(\text{Grad rot } v)^\top$
- $2 \text{spn}^{-1} \text{skw Rot } S = \text{Div } S^\top - \text{grad tr } S = \text{Div}(S - (\text{tr } S) \text{ id})^\top$ , in particular,  $\text{rot Div } S^\top = 2 \text{rot spn}^{-1} \text{skw Rot } S$  and  $2 \text{skw Rot } S = \text{spn Div } S^\top$ , if  $\text{tr } S = 0$
- $\text{tr Rot } S = 2 \text{div spn}^{-1} \text{skw } S$ , in particular,  $\text{tr Rot } S = 0$ , if  $\text{skw } S = 0$ , and  $\text{tr Rotsym } S = 0$  and  $\text{tr Rot skw } S = \text{tr Rot } S$
- $2(\text{Grad spn}^{-1} \text{skw } S)^\top = (\text{tr Rot skw } S) \text{ id} - 2 \text{Rot skw } S$
- $3 \text{Div}(\text{dev Grad } v)^\top = 2 \text{grad div } v$
- $2 \text{Rot sym Grad } v = -2 \text{Rot skw Grad } v = -\text{Rot spn rot } v = (\text{Grad rot } v)^\top$

- $2\text{Divsym Rot}S = -2\text{Divskw Rot}S = \text{rot Div}S^\top$
- $\text{Rot}(\text{Rotsym}S)^\top = \text{sym Rot}(\text{Rot}S)^\top$
- $\text{Rot}(\text{Rotskw}S)^\top = \text{skw Rot}(\text{Rot}S)^\top$

All formulas extend also to distributions.

## APPENDIX B: ELASTICITY COMPLEX OPERATORS REVISITED

Let  $\top$  denote the formal operator of matrix transposition, that is,

$$\top S := S^\top,$$

and define

$$\text{axl} : \mathbb{R}_{\text{skw}}^{3 \times 3} \rightarrow \mathbb{R}^3; \begin{bmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{bmatrix} \mapsto \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix}.$$

We recall the operators forming the de Rham complex (classical vector analysis) grad, rot, and div acting on functions and vector fields, respectively, as formal matrix operators

$$\text{grad} := \begin{bmatrix} \partial_1 \\ \partial_2 \\ \partial_3 \end{bmatrix}, \text{rot} := \text{axl}^{-1} \text{grad} = \begin{bmatrix} 0 & -\partial_3 & \partial_2 \\ \partial_3 & 0 & -\partial_1 \\ -\partial_2 & \partial_1 & 0 \end{bmatrix}, \text{div} := \top \text{grad} = [\partial_1 \ \partial_2 \ \partial_3].$$

Moreover, we introduce their relatives from the vector de Rham complex acting on vector and tensor fields, respectively, as formal matrix operators

$$\text{Grad} := \top \text{grad} \top, \quad \text{Rot} := \top \text{rot} \top, \quad \text{Div} := \top \text{div} \top.$$

In words, Grad, Rot, and Div act row-wise as the operators grad, rot, and div from the classical de Rham complex. Note that in previous works,<sup>8–11</sup> we used the notation  $\text{spn} = \text{axl}^{-1}$  and that  $\text{Grad}v$  is just the Jacobian for a vector field  $v$ .

Let

$$i_{\mathbb{S}} : \mathbb{R}_{\text{sym}}^{3 \times 3} \rightarrow \mathbb{R}^{3 \times 3}$$

denote the canonical embedding of symmetric  $(3 \times 3)$ -matrices into the arbitrary  $(3 \times 3)$ -matrices. Then the adjoint

$$i_{\mathbb{S}}^* : \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}_{\text{sym}}^{3 \times 3}$$

is almost the projector onto symmetric  $(3 \times 3)$ -matrices, that is, the actual projector is given by

$$\text{sym} := i_{\mathbb{S}}^* i_{\mathbb{S}} : \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}_{\text{sym}}^{3 \times 3}; S \mapsto \frac{1}{2}(S + S^\top).$$

We extend  $\top$ ,  $\text{axl}$ ,  $\text{axl}^{-1}$ ,  $i_{\mathbb{S}}$ ,  $i_{\mathbb{S}}^*$ , and  $\text{sym}$  to  $L^2(\Omega)$ -tensor fields.

In the light of this, in symmetric elasticity theory, we are dealing with the operators

$${}_{\mathbb{S}}\text{Grad} := i_{\mathbb{S}}^* \text{Grad}, \quad {}_{\mathbb{S}}\text{Rot} \top \text{Rot}_{\mathbb{S}} := i_{\mathbb{S}}^* \text{Rot} \top \text{Rot}_{\mathbb{S}} = i_{\mathbb{S}}^* \top \text{rot} \top \text{rot} \top i_{\mathbb{S}}, \quad \text{Div}_{\mathbb{S}} := \text{Div}_{i_{\mathbb{S}}},$$

and their maximal  $L^2(\Omega)$ -realizations, that is,

$$\begin{aligned} {}_{\mathbb{S}}\text{Grad} &: D({}_{\mathbb{S}}\text{Grad}) \subset L^2(\Omega) \rightarrow L^2_{\mathbb{S}}(\Omega), \\ {}_{\mathbb{S}}\text{Rot} \top \text{Rot}_{\mathbb{S}} &: D({}_{\mathbb{S}}\text{Rot} \top \text{Rot}_{\mathbb{S}}) \subset L^2_{\mathbb{S}}(\Omega) \rightarrow L^2_{\mathbb{S}}(\Omega), \\ \text{Div}_{\mathbb{S}} &: D(\text{Div}_{\mathbb{S}}) \subset L^2_{\mathbb{S}}(\Omega) \rightarrow L^2(\Omega). \end{aligned}$$

Note that  $\iota_{\mathbb{S}} \mathbb{S} \text{Grad} = \text{symGrad}$  and  $\iota_{\mathbb{S}} \mathbb{S} \text{Rot} \top \text{Rot}_{\mathbb{S}} = \text{symRot} \top \text{Rot}_{\mathbb{S}}$ . Moreover, on symmetric tensor fields, we have  $\mathbb{S} \text{Rot} \top \text{Rot}_{\mathbb{S}} = \text{Rot} \top \text{Rot}$  (cf. lemma A.1<sup>10,11</sup>)

Finally, the operators  $\text{symGrad}_{\Gamma_i}$ ,  $\text{RotRot}_{\mathbb{S}, \Gamma_i}^{\top}$ , and  $\text{Div}_{\mathbb{S}, \Gamma_i}$  from Section 2.2 are the restrictions of  $\mathbb{S} \text{Grad}$ ,  $\mathbb{S} \text{Rot} \top \text{Rot}_{\mathbb{S}}$ , and  $\text{Div}_{\mathbb{S}}$  to the domains  $D(\text{symGrad}_{\Gamma_i})$ ,  $D(\text{RotRot}_{\mathbb{S}, \Gamma_i}^{\top})$ , and  $D(\text{Div}_{\mathbb{S}, \Gamma_i})$ , which are the closures of  $C_{\Gamma_i}^{\infty}(\Omega)$  and  $C_{\mathbb{S}, \Gamma_i}^{\infty}(\Omega)$  in the corresponding graph norms, respectively.